

Dynamics of surface diffeomorphisms relative to homoclinic and heteroclinic orbits

Pieter Collins *

Department of Mathematical Sciences

University of Liverpool

Liverpool L69 7ZL, U.K.

pieter.collins@12move.nl

February 1, 2008

Abstract

In this paper, we show how to obtain information about the dynamics of a two-dimensional discrete-time system from its homoclinic and heteroclinic orbits. The results obtained are based on the theory of *trellises*, which comprise finite-length subsets of the stable and unstable manifolds of a collection of saddle periodic orbits. For any collection of homoclinic or heteroclinic orbits, we show how to associate a canonical *trellis type* which describes the orbits. Given a trellis type, we show how to compute a *graph representative* which gives a combinatorial invariant of the trellis type. The orbits of the graph representative represent the dynamics forced by the homoclinic/heteroclinic orbits in the sense that every orbit of the graph representative is *globally shadowed* by some orbit of the system, and periodic, homoclinic/heteroclinic orbits of the graph representative are shadowed by similar orbits. By constructing suitable surface diffeomorphisms, we prove that these results are optimal in the sense that the topological entropy of the graph representative is the infimum of the topological entropies in the isotopy class relative to the homoclinic/heteroclinic orbits.

Mathematics subject classification: Primary: 37E30. Secondary: 37B10, 37C27, 37E25.

*This work was partially funded by Leverhulme Special Research Fellowship SRF/4/9900172

Contents

1	Introduction	3
2	Tangles and Trellises	6
2.1	Tangles	6
2.2	Trellises	7
2.3	Attractors	12
2.4	Extensions	14
2.5	Reducibility	15
2.6	Cutting	19
2.7	Curves	19
3	Biasymptotic Orbits and Minimal Trellises	22
3.1	Biasymptotic orbits	22
3.2	Existence of minimal trellises	25
3.3	Uniqueness of minimal trellis	29
3.4	Minimal extensions	31
4	Graphs Representatives	32
4.1	Combinatorics of graph maps	32
4.2	Compatible graph maps	34
4.3	Properties of graph representatives	38
4.4	Reducibility of graph representatives	41
4.5	Uniqueness of graph representatives	43
5	Computing a Graph Representative	44
5.1	Graph moves	45
5.2	Effect of graph moves	47
5.3	Algorithm to compute the graph representative	49
5.4	Algorithm for reduction	54
6	Symbolic Dynamics and Shadowing	55
6.1	Symbolic dynamics	55
6.2	Global shadowing	56
6.3	Shadowing and Nielsen numbers	58
6.4	Shadowing by trellis orbits	59
6.5	Shadowing by stable and biasymptotic orbits	61
7	Entropy-Minimising Diffeomorphisms	63
7.1	Nielsen entropy of minimal trellises	65
7.2	Regular domains	69
7.3	Existence of entropy minimisers	75
7.4	Approximate entropy minimising diffeomorphisms	76
7.5	Non-existence of entropy minimisers	81
A	Intersections and Isotopies of Curves in Surfaces	82

A.1	Minimal intersections	83
A.2	Ambient isotopies of curves	84
A.3	Nielsen intersection theory	85

1 Introduction

The importance of homoclinic orbits in dynamical systems theory was first realised by Poincaré [Poi99], who showed that the presence of a homoclinic tangle in the three-body problem was enough to show that the system was non-integrable. More work on homoclinic tangles was later undertaken by Birkhoff [Bir49], but modern interest was inspired by Smale, who showed that any system with a transverse homoclinic point must have a horseshoe in some iterate [Sma63]. In this paper, we extend the local analysis of Smale to obtain a global understanding of the dynamics for diffeomorphisms in two dimensions. The analysis depends on constructing a *graph representative* of the system, from which we obtain information about the dynamics in terms of *symbolic dynamics* relative to a Markov partition and *global shadowing*.

The use of Markov partitions to give symbolic dynamics is a fundamental tool in dynamical systems theory. For uniformly-hyperbolic systems, the dynamics is described completely by the symbolics, but for non-uniformly hyperbolic systems, the Markov partition typically only gives a *lower bound* for the dynamics. From this, we can often deduce information on the *forcing relation*, which describes when the presence of a particular geometric structure implies the existence of another structure. Sharkovskii's theorem for interval maps is a classic example of a forcing result: a total ordering $\succ$ of the natural numbers is given such that the presence of a periodic orbit of period p forces periodic orbits of all periods q with $p \succ q$. Sharkovskii's theorem can be strengthened to consider the ordering of points on the orbit, and this gives a complete (though less elegant) description of the forcing relation for periodic orbits of interval maps. Interval maps can be studied by elementary methods, but obtaining symbolic dynamics for more complicated classes of system typically requires the use of some topological index theory, such as the fixed point index [Fri83, Jia93] or the Conley index [Con78].

The theory of isotopy classes of surface diffeomorphisms relative to periodic orbits is known as *Nielsen-Thurston theory* [CB88]. Here, the period is no longer a useful characterisation of the orbit; instead, we need to consider the *braid type* which is the conjugacy class of isotopy classes relative to the orbit in question. Just as for periodic orbits of interval maps, orbits of different periods may have different braid types. For each braid type there is a canonical representative such that a braid type BT_1 forces BT_2 if and only if the *Thurston minimal representative* of BT_1 contains an orbit of type BT_2 . A Thurston minimal representative has the minimum number of periodic points of any period, and minimal topological entropy in the isotopy class [Han85]. The most interesting case is when the canonical representative is pseudo-Anosov. A pseudo-Anosov map preserves a pair of transverse foliations, the leaves of the stable foliation being uniformly contracted and the unstable foliation being uniformly expanded. Pseudo-Anosov maps therefore exhibit a type of uniformly hyperbolic behaviour, and Thurston's classification theorem

can be seen as a persistence result for this hyperbolicity.

Unfortunately, Thurston's original proof was non-constructive, and thus of little use in actually determining the forcing relation. It was not until an algorithm of Los [Los93] that computations became possible. Los' algorithm was soon followed by an algorithm which was discovered independently by Franks and Misiurewicz [FM93] and Bestvina and Handel [BH95]. A description of the moves of this algorithm with an emphasis on implementation details was given by Keil [Kei97]. Even so, the global properties of the forcing relation are still only partially understood and this is an active area of research [dCH].

A more challenging problem is to consider the dynamics of surface diffeomorphisms relative to homoclinic and heteroclinic orbits of tangles. The standard results of Poincaré and Smale for transverse homoclinic points have since been extended to topologically transverse intersections which are not geometrically transverse [BW95, Ray98], and a great deal is known about bifurcations near a homoclinic tangency [New80, MV93]. The global dynamics and the forcing relation have been previously considered by McRobie and Thompson [MT94], who carried out a detailed study of the *spiral 3-shoe*, by Rom-Kedar [RK94], who obtained results on entropy bounds and homoclinic forcing using a *topological approximation method*, and by Easton [Eas98]. Algorithms for computing dynamics forced by homoclinic orbits were recently given independently by Handel [Han99] and Hulme [Hul00]. Homoclinic tangles have been used by Christiansen and Politi [CP96] to compute generating partitions for symbolic dynamics. Orbits of the Hénon map have also been obtained using continuation methods by Sterling, Dullin and Meiss [SDM99].

The aim of this paper is to give a complete and general theoretical basis for describing and computing the dynamics forced by a homoclinic or heteroclinic tangle. The results are a completion of the theory begun in [Col99], and are related to the pruning theory of de Carvalho [dC99]. Instead of considering an entire tangle, which consists of immersed curves of infinite length, we only consider a subset consisting of curves of finite length. We call such a figure a *trellis*, after Birkhoff [Bir49] and Easton [Eas86]. Aside from being easier to work with, trellises have the advantage that there exist a number of algorithms to compute them to a very high degree of accuracy, such as those of Simo [Sim89] and Krauskopf and Osinga [KO98]. We therefore obtain results which can be directly applied to physically-relevant systems. In Section 2 we give a formal description of trellises and discuss the features of their geometry which we shall need later.

In Section 3, we show how to relate a collection of homoclinic and/or heteroclinic orbits to a canonical trellis type forcing the same dynamics, thus unifying the viewpoints of homoclinic/heteroclinic orbits and tangles. This result can be applied to classify homoclinic orbits by braid type. In particular, it is straightforward to obtain a classification of *horseshoe homoclinic orbits*, which are homoclinic orbits of the Smale horseshoe map.

For the rest of the paper, we try to mimic the results of Nielsen-Thurston theory as closely as possible. While the fact that we are dealing with homoclinic orbits often complicates matters, it sometimes can yield simplifications. In Section 4, we show how to associate a graph representative to a given trellis map, in much the same way that a pseudo-Anosov braid type gives rise to a train track. The graph we obtain is unique (unlike a train track)

and gives a convenient way of specifying the trellis. From this, we also obtain a canonical graph representative for a homoclinic orbit.

We consider the problem of computing the graph representative in Section 5. We give an algorithm modelled on the Bestvina-Handel algorithm, but again the invariant manifolds simplify matters by enabling us to avoid the notorious valence-2 homotopies of the graph map. We follow an idea of Los [Los96] by replacing the valence-2 homotopy by a *valence-3 homotopy* which does not require a global knowledge of the map. Unfortunately, Los' analysis of the effect of the move on entropy was erroneous; the entropy may increase during this move, and this meant that the algorithm was not proved to terminate. Our analysis goes along different lines and avoids entropy computations altogether. We also unify the process of absorbing into the peripheral subgraph with the main algorithm, which also simplifies matters somewhat. As with the Bestvina-Handel algorithm, reductions are obtained and dealt with accordingly. Finally, we give a simplified algorithm for braid types based on our analysis of the algorithm for trellises.

In Section 6, we discuss how the dynamics of the graph representative model the dynamics of the surface diffeomorphism. These results are based on the *relative periodic point theory* developed in [Col01]. We first deduce the existence of periodic orbits; the existence of other orbits follows by taking limits. In particular, these results show that the topological entropy of the graph representative is a lower bound for the topological entropy of the surface diffeomorphism. This lower bound is called the *Nielsen entropy*, since it is related to the growth rate of the number of essential Nielsen classes of period- n orbits. Orbits are related by *global shadowing*, a relation introduced by Katok, though here we give a new definition more in keeping with the spirit of our topological analysis. One important fact is that periodic orbits which globally shadow each other have the same braid type, so we obtain information about the braid types of the orbits of the system. Finally, we show by a careful construction that homoclinic and heteroclinic orbits of the graph map are shadowed by homoclinic and heteroclinic orbits of the surface diffeomorphism.

The final section, Section 7, is devoted to showing that the dynamics computed by our algorithm is indeed all the dynamics forced by the trellis. In particular, the lower entropy bound computed by the algorithm is sharp. Unfortunately, unlike Nielsen-Thurston theory, there may not be a canonical diffeomorphism whose topological entropy equals the entropy bound given by the Nielsen entropy. We give necessary and sufficient conditions for the existence of a diffeomorphism whose entropy equals the Nielsen entropy. We also show that by carefully extending the original trellis, we can always find a diffeomorphism whose entropy approximates the Nielsen entropy arbitrarily closely. Finally, we complete the link with Nielsen-Thurston theory by finding pseudo-Anosov maps which approximate the Nielsen entropy when such maps exist for the given trellis type.

A unifying theme throughout this paper is to consider isotopy classes of curves embedded in the surface. We are particularly concerned with curves that have the minimum possible number of intersections with the stable curves of the trellis. Stable curves can be thought of as cohomology classes, though we do not use this interpretation here, as the abelian homology theory is insufficient for many of our constructions, which require the non-abelian homotopy groups. Many of the results we need are intuitively obvious “folk theorems,” but we give complete proofs in Appendix A.

2 Tangles and Trellises

In this section we discuss the geometry of homoclinic and heteroclinic tangles, and introduce trellises, trellis mapping classes and trellis types, which are our main topic of study. We give a number of definitions which allow us to describe some of the properties of trellises. We discuss reducibility of trellis mapping classes, and explain how this differs from reducibility of mapping classes of surface diffeomorphisms. Finally, we discuss the procedure of cutting along the unstable set, which gives us a way of relating trellis maps to maps of topological pairs, and hence a way of studying the dynamics.

2.1 Tangles

A tangle is the figure formed by the stable and unstable manifolds of a collection of periodic saddle orbits.

Definition 2.1 (Tangle) Let f be a diffeomorphism of a surface M , and P a finite invariant set of saddle points of f . Let $W^S(f; P)$ be the stable set of P , and $W^U(f; P)$ the unstable set of P . The pair $W = (W^U(f; P), W^S(f; P))$ is a *tangle* for f . If P consists of a single point, then W is a *homoclinic* tangle, otherwise W is a *heteroclinic* tangle.

If p is a point of P , then the stable and unstable curves passing through p are denoted $W^S(f; p)$ and $W^U(f; p)$ respectively.

We use the notation $W^{U/S}$ in statements which hold for both stable and unstable manifolds. If the diffeomorphism f under consideration is clear from the context, we abbreviate $W^{U/S}(f; P)$ to $W^{U/S}(P)$, and if the periodic point set P is also clear, we simply write $W^{U/S}$.

We use the obvious notation for intervals of $W^{U/S}$: The closed interval in $W^{U/S}$ with endpoints a and b is denoted $W^{U/S}[a, b]$ and the open interval is denoted $W^{U/S}(a, b)$. Half-open intervals are denoted $W^{U/S}[a, b)$ and $W^{U/S}(a, b]$.

Each stable or unstable curve $W^{U/S}(p)$ is naturally separated into two pieces by the periodic point p .

Definition 2.2 (Branch) A *branch* of a tangle is a component of $W^{U/S}(P) \setminus P$. We orient each branch of $W^S(p)$ so that the positive orientation is in the direction of p , and each branch of $W^U(p)$ so that the positive orientation is away from p . A branch is *trivial* if it does not intersect any other branch.

We denote a branch of $W^{U/S}(p)$ by integers $W^{U/S}(p, b)$. Note that the branches of $W^{U/S}$ are permuted by f .

Definition 2.3 (Fundamental domain) A *fundamental domain* is a half-open interval in $W^{U/S}$ bounded by points q and $f^n(q)$, where $q \notin W^P$, and n is the least integer such that q and $f^n(q)$ lie in the same branch.

A branch of a tangle can be found by taking the union of the n -th iterates of a fundamental domain.

Definition 2.4 (Biasymptotic points) An *intersection point* of a tangle $W(P)$ is a point in $W^U \cap W^S$. An intersection point not in P is a *biasymptotic* point; an *asymptotic* point is a point in $(W^U \cup W^S) \setminus P$. A biasymptotic point $q \in W^U(p_1) \cap W^S(p_2)$ is a *homoclinic* point if p_1 and p_2 are part of the same periodic orbit, otherwise q is a *heteroclinic* point. If W^U and W^S cross transversely at a biasymptotic point q , then q is a *transverse* homoclinic or heteroclinic point.

There are six different types of isolated intersection point, two transverse intersections $\times$, with positive (+) and negative (−) crossings respectively, and are four tangential intersections $\succ$. It is well known [Sma63] that if a diffeomorphism f has a transverse homoclinic point, then f must be chaotic and have positive topological entropy. If f is a uniformly hyperbolic diffeomorphism, then any biasymptotic point is a transverse intersection.

Definition 2.5 (Primary intersection point) An intersection point $q \in W^U(p_1) \cap W^U(p_2)$ is a *primary intersection point* or *pip* if $W^U(p_1, q)$ and $W^S(q, p_2)$ are disjoint.

2.2 Trellises

There are a number of difficulties in working with tangles directly. These are due to the fact that if any branch intersects another, then it will be an immersed, rather than embedded, curve. The geometry of such a curve is extremely complicated, as is the global topology of the tangle. Further, since branches may have infinite length, tangles are impossible to compute directly. Instead, we work with subsets of tangles known as *trellises*, for which the stable and unstable curves have finite lengths.

Definition 2.6 (Trellis) Let f be a diffeomorphism of a surface M with a finite invariant set P of hyperbolic saddle points. A *trellis* for f is a pair $T = (T^U, T^S)$, where T^U and T^S be subsets of $W^U(f; P)$ and $W^S(f; P)$ respectively such that

1. T^U and T^S both consist of finitely many compact intervals with non-empty interiors, and
2. $f(T^U) \supset T^U$ and $f(T^S) \subset T^S$.

We denote the set of periodic points of T by T^P , and the set of intersections of T^U and T^S by T^V .

Definition 2.7 (Trellis map) If f is a diffeomorphism and T is a trellis for f , then the pair $(f; T)$ is called a *trellis map*.

There are two important relations between trellis maps, namely *isotopy* and *conjugacy*.

Definition 2.8 (Isotopy and conjugacy) Trellis maps $(f_0; T)$ and $(f_1; T)$ are *isotopic* if there is an isotopy of surface diffeomorphisms (f_t) such that for each t , $(f_t; T)$ is a trellis

map. Trellis maps $(f_0; T_0)$ and $(f_1; T_1)$ are *conjugate* if there is a homeomorphism h such that $h \circ f_0 = f_1 \circ h$ and $h(T_0^{U/S}) = T_1^{U/S}$.

These these relations allow us to define equivalence classes of trellis map. It is these equivalence classes which will be our primary object of study.

Definition 2.9 (Trellis mapping class and trellis type) The *trellis mapping class* $([f]; T)$ is the set of all trellis maps which are isotopic to f . The *trellis type* $[f; T]$ is the set of all trellis maps which are conjugate to a map in $([f]; T)$. Trellis maps $(f_0; T_0)$ and $(f_1; T_1)$ are considered equivalent if $[f_0; T_0] = [f_1; T_1]$.




Figure 1: Trellises in the Hénon map. (a) is a homoclinic trellis, (b) a heteroclinic trellis.

In Figure 1 we show two trellises, each of which is homeomorphic to a trellis for a diffeomorphism in the Hénon family. In each case the small crosses mark primary intersection points on a single orbit. It turns out that this is enough information to specify a trellis type for these trellises.

Just as we can define branches of a tangle, we can also define branches of a trellis.

Definition 2.10 (Branch) If $W^{U/S}(p, b_{u/s})$ is branch of a tangle for f , then the corresponding branch of a trellis $T \subset W$ is given by

$$T^{U/S}(p, b_{u/s}) = \{p\}\{p\} \cup W^{U/S}(p, b_{u/s}) \cap T^{U/S}.$$

Note that a branch of a trellis is a union of embedded closed intervals, whereas a branch of a tangle is an immersed open interval. In particular, a branch of a trellis contains the periodic saddle point.

Most numerical algorithms compute trellises by growing out the branches from P , and the resulting trellises have connected branches. While this is the most natural case to visualise, it is often important to consider trellises with branches that are not connected. Disconnected branches can be useful when considering trellises which are subsets of tangles with unbounded branches.

The most important subsets of $T^{U/S}$ are intervals with endpoints in T^V , especially those which have no intersections in their interior.

Definition 2.11 (Arc, segment and end) An *arc* of $T^{U/S}$ is a closed subinterval of $T^{U/S}$ with endpoints in T^V . A *segment* of $T^{U/S}$ is an arc of $T^{U/S}$ with no topologically transverse intersection points in its interior. An *end* of a trellis is a subintervals of $T^{U/S}$ which does not lie in any segment. An *end intersection* of a trellis is an intersection which does not lie in two segments of $T^{U/S}$ (so lies in at least one end of $T^{U/S}$).

Similarly, the most important subsets of the surface M are those bounded by T^U and T^S .

Definition 2.12 (Domain and region) An *open domain* of T is an open subset of M with boundary in $T^U \cup T^S$. A *closed domain* is the closure of an open domain. An *open region* of T is a component of $M \setminus (T^U \cup T^S)$. A *closed region* is the closure of an open region, and hence includes the stable and unstable boundary segments. We will denote domains by the letter D , and regions by the letter R .

There are two special types of region which play an important role later.

Definition 2.13 (Rectangle) A *rectangle* is a region which is a topological disc bounded by two stable and two unstable segments which make acute or obtuse angles with each other (i.e. do not form reflex angles).

Definition 2.14 (Bigon) A *bigon* is a region which is a topological disc bounded by one stable and one unstable segment which make acute or obtuse angles with each other.

Bigons play a similar role in the trellis theory as do critical points in the kneading theory, and may be called *critical* regions.

In general, we do not have much control over the geometry of a trellis. The exception is near a point of T^P , where the dynamics are conjugate to a linear map. It is important to consider how the nontrivial branches T divide the surface in a neighbourhood of T^P .

Definition 2.15 (Quadrant, secant and coquadrant) Let T be a trellis, and $\bar{T}$ be the nontrivial branches of T , and p be a point of T^P . Then a *sector* of T at p is a local component of a region of $\bar{T}$ in a neighbourhood of p . A sector is a *quadrant* Q if it intersects no trivial branches, in which case the boundary includes a single unstable branch $T^U(Q)$ and a single stable branch $T^S(Q)$. A sector is an *secant* if it intersects a single trivial branch; if this is a branch of T^U the secant is *attracting*, and if this is a branch of T^S , the secant is repelling. A sector is a *coquadrant* if it intersects two trivial branches, in which case the only other sector at p is a quadrant.

Quadrants, secants and coquadrants are shown in Figure 2. The region containing a quadrant Q is denoted $R(Q)$. The *image* of a quadrant Q is the quadrant containing $f(Q)$; note that $T^{U/S}(f(Q))$ is the same branch as $f(T^{U/S}(Q))$. A stable segment S with endpoint $q \in T^U(Q)$ intersects lies on the Q -side of T^U if locally S lies on the same side of $T^U(Q)$ at q as $T^S(Q)$ does at p . Similarly, an unstable segment U with endpoint $q \in T^S(Q)$ intersects lies on the Q -side of T^S if locally U lies on the same side of $T^S(Q)$ at q as $T^U(Q)$ does at p . The segment S in Figure 2(a) lies on the Q -side of q .




Figure 2: Sectors at p . (a) is a quadrant, (b) an attracting secant, (c) a repelling secant and (d) a coquadrant.

We now illustrate the concepts introduced so far with one of the most important trellis types, the *Smale horseshoe trellis*.




Figure 3: The Smale horseshoe trellis.

Example (The Smale horseshoe trellis)

The Smale horseshoe trellis is formed by the stable and unstable manifolds of the direct saddle fixed point (i.e. the saddle point with positive eigenvalues) of the Smale horseshoe map, and is depicted in Figure 3. The stable and unstable sets are subsets of the stable and unstable manifolds of the saddle fixed point p . The branches of this trellis are connected and the all intersection points are transverse. The points q_0 , q_1 and q_2 are primary intersection points on a single homoclinic orbit. The orbits of the biasymptotic points v_0 and v_1 are shown in white dots. These orbits are called *forcing orbits* for the Smale horseshoe trellis; it is essentially impossible to remove any intersection points of the Smale horseshoe trellis by an isotopy without first removing v_0 and v_1 in a homoclinic bifurcation.

Throughout this paper we follow the convention of showing forcing orbits as which dots, and primary intersection points as crosses.

There are eight regions, an unbounded region R_∞ , three bigons and four rectangles. The quadrant Q is contained in the region R_0 . Under the Smale horseshoe map, there is a Cantor set of nonwandering points contained in the (closed) rectangular regions R_0 and R_1 , including a fixed point b in R_1 . All other points are wandering except for an attracting fixed point at the end of one unstable branch of $T^U(p)$ in R_∞ . The topological entropy of the Smale horseshoe map is $\log 2$. We shall see later that any diffeomorphism with this trellis type must have topological entropy $h_{top} \geq \log 2$.

Notice that one of the unstable branches of the Smale horseshoe trellis ends in an attracting fixed point, and one of the stable branches is non-existent. Both of these are therefore trivial branches.




Figure 4: The singular Smale horseshoe trellis.

We could isotope the Smale horseshoe map in a neighbourhood of the saddle point p to obtain a homeomorphism with a one-prong singularity at p . The new trellis is shown in Figure 4. It is topologically equivalent to the Smale horseshoe trellis (minus the unstable branch ending at a) but has a different geometric type, and is known as the *singular Smale horseshoe trellis*. Many of the properties of the singular Smale horseshoe trellis are the same as that of regular Smale horseshoe trellis, except that now there is no homeomorphism in the trellis mapping class with a one-prong singularity at p which has topological entropy $\log 2$. Instead, for any $\epsilon > 0$ there is a pseudo-Anosov homeomorphism f with topological entropy $\log 2 < h_{top}(f) < \log 2 + \epsilon$.

The most important dynamical feature of a trellis type is its *entropy*.

Definition 2.16 (Entropy) The *entropy* of a trellis type $[f; T]$, denoted $h_{top}[f; T]$ is the infimum of the topological entropies of diffeomorphisms in $[f; T]$; that is

$$h_{top}[f; T] = \inf\{h_{top}(\hat{f}) : \hat{f} \in [f; T]\}.$$

If this infimum is not a minimum, we sometimes write $h_{top}[f; T] = \inf\{h_{top}(\hat{f})\} + \epsilon$.

We finally give two subclasses of trellis maps which are important for technical reasons. In both these subclasses, we treat the stable and unstable sets differently; this anticipates the loss of time-reversal in the cutting procedure of Section 2.6. We consider *proper trellises* in order that the stable set be well-behaved after cutting.

Definition 2.17 (Proper trellis) We say a trellis $T = (T^U, T^S)$ is *proper* if $\partial T^S \subset T^U$ and $\partial T^U \cap T^S = \emptyset$. In other words, a trellis is proper if the endpoints of intervals in T^S lie in T^U , but the endpoints of intervals in T^U do not lie in T^S . A trellis map $(f; T)$ is proper if $f^n(U)$ does not contain a point of T^V for any open unstable end U (so the ends of T^U do not force any dynamics).

Note that considering proper trellises is no great loss of generality; an unstable curve with an endpoint in T^S can be extended by an arbitrarily short piece of curve, and a stable

curve with an endpoint not in T^U can be shrunk by removing its end. Further, since the ends of a trellis give no information about the dynamics, both these procedures are harmless and do not affect dynamical computations in any way. Henceforth, all trellises will be taken to be proper trellises unless otherwise stated.

We consider *well-formed* trellises in order that the isotopy classes of curves with endpoints in T^S are well behaved under iteration. This will be important when considering the graph representative defined in Section 4.2, as it gives a necessary condition for the topological entropy of the graph representative to be an optimal entropy bound for the trellis mapping class.

Definition 2.18 (Well-formed trellis) We say a proper trellis $T = (T^U, T^S)$ for a diffeomorphism f is *well-formed* if every component of $T^U \cup f(T^S)$ contains a point of T^P .

Again, taking well-formed trellises is no great loss of generality, as we will usually take trellises for which the unstable branches are connected, and it remains to ensure that for every stable segment interval S , its image $f(S)$ intersects T^U , which we can do by removing the stable intervals whose iterates do not intersect T^U .

2.3 Attractors

As well as saddle periodic orbits, generic diffeomorphisms also have attracting and repelling periodic orbits. Clearly, a stable segment cannot be in the basin of attraction of an attracting orbit, and neither can an unstable segment be in the basin of a repelling orbit. However, it is possible for the interior of an entire region to lie in the basin of an attracting or repelling periodic orbit. Such regions are called *stable* or *unstable* regions.

Definition 2.19 (Stable and unstable regions) Let $([f]; T)$ be a trellis mapping class. A domain D of T is *stable* if there is a diffeomorphism $\tilde{f}$ in $([f]; T)$ and a periodic orbit P of f such that $\tilde{f}^n(x) \rightarrow P$ as $n \rightarrow \infty$ for all x in the interior of D . If D contains a point of P , then D is *attracting*.

Similarly, a domain D is *unstable* if there is a diffeomorphism $\tilde{f}$ in $([f]; T)$ and a periodic orbit P of f such that $\tilde{f}^{-n}(x) \rightarrow P$ as $n \rightarrow \infty$ for all x in the interior of D , and if D contains a point of P , then D is *repelling*.

Two examples of attracting domains are shown in Figure 5. All points in the shaded domains are in the basin of attraction of the stable periodic point. Note that T^S is disjoint from the interior of A . Repelling domains are similar. Stable and unstable domains are wandering sets, unless they contain the attracting or repelling periodic orbit.

A region which contains an attractor and a repeller is called *unbounded* region, by analogy with the unbounded region of the planar Smale horseshoe, for which the repeller is the point at infinity. Figure 6 shows an unbounded region R near a fixed point p . All points in A lie in the basin of attraction of an attracting fixed point a , and all points in B lie in




Figure 5: Attracting domains.




Figure 6: An unbounded region.

the basin of a repelling fixed point b . Since A and B cover the interior of R , all points in the interior of R are wandering, apart from a and b .

An open stable segment which is in the basin of a repelling periodic orbit is also a wandering set, as is an open unstable segment which is in the basic of an attracting periodic orbit. There are some nonwandering segments such that any finite iterate need not contain an intersection point. We call such segments *almost wandering*.

Definition 2.20 (Almost wandering segment) Let $([f]; T)$ be a trellis mapping class. An open segment is *almost wandering* if it is nonwandering, but for any integer n there exists a diffeomorphism $\tilde{f} \in ([f]; T)$ the n^{th} iterate of the segment by $\tilde{f}$ contains no intersection points.

Therefore, an open stable segment S is almost wandering if for any positive integer n , there exists $\tilde{f} \in ([f]; T)$ such that $\tilde{f}^{-n}(S) \cap T^U = \emptyset$, but for any diffeomorphism $\hat{f} \in ([f]; T)$, there exists n such that $\hat{f}^{-n}(S) \cap T^U \neq \emptyset$. A similar statement holds for unstable segments.

Definition 2.21 (Chaotic region) Let $([f]; T)$ be a trellis mapping class. A region R is *chaotic* if for every diffeomorphism $\hat{f}$ in $([f]; T)$, there exists an integer n such that $\hat{f}^n(T^U) \cap \hat{f}^{-n}(T^S)$ contains a point in the interior of R .

A chaotic region supports entropy, in the sense that for any diffeomorphism $\hat{f} \in ([f]; T)$, there is an ergodic measure μ with positive entropy such that $\mu(R) > 0$.

2.4 Extensions

A surface diffeomorphism f with a periodic saddle orbit has infinitely many trellises, which are partially ordered by inclusion. Taking a smaller trellis gives a *subtrellis*, and a larger trellis a *supertrellis*.

Definition 2.22 (Subtrellis and supertrellis) Let $(f; T)$ and $(f; \hat{T})$ be trellis maps. If $T^{U/S} \subset \hat{T}^{U/S}$, then T is a *subtrellis* of $\hat{T}$, and $\hat{T}$ is a *supertrellis* of T . Similarly, if $([f]; T)$ is a trellis mapping class, a trellis map $(\hat{f}; \hat{T})$ is a *supertrellis* for $([f]; T)$ if $\hat{f} \in ([f]; T)$ and $T^{U/S} \subset \hat{T}^{U/S}$.

Since taking a subtrellis loses information about the trellis map, we are more interested in supertrellises. Of particular importance are those trellises which can be obtained by iterating segments or branches.

Definition 2.23 (Iterate) Let $(f; T)$ be a trellis map. Then a trellis $\hat{T}$ is an *f-iterate* of f if there exist positive integers n_u and n_s such that $\hat{T}^U = f^{n_u}(T^U)$ and $\hat{T}^S = f^{-n_s}(T^S)$. An iterate is a *stable iterate* if $\hat{T}^U = T^U$ and an *unstable iterate* if $\hat{T}^S = T^S$. A trellis map $(\hat{f}; \hat{T})$ is an iterate of the trellis mapping class $([f]; T)$ if $\hat{f} \in ([f]; T)$ and $\hat{T}$ is an $\hat{f}$ -iterate of T .

The concept of an iterate is rather restrictive in practice, as we may wish to iterate only a few segments of T . Further, trellises are often computed by growing branches out from T^P , so we may obtain a supertrellis for which $\hat{T}^P = T^P$ but which is not an iterate. A class of supertrellis which is more restrictive than that of an iterate is an *extension*.

Definition 2.24 (Extension) Let $(f; T)$ be a trellis map. A trellis $\hat{T}$ is an *f-extension* of T if there exists n such that

$$T^U \subset \hat{T}^U \subset f^n(T^U) \quad \text{and} \quad T^S \subset \hat{T}^S \subset f^{-n}(T^S).$$

A trellis map $(\hat{f}; \hat{T})$ is an *extension* of a trellis mapping class $([f]; T)$ if $\hat{f} \in ([f]; T)$ and $\hat{T}$ is a $\hat{f}$ -extension of T .

In other words, a trellis $\hat{T}$ is an extension of T if it is a subtrellis of some iterate of T . Just as for iterates, a *stable extension* has $\hat{T}^U = T^U$ and an *unstable extension* has $\hat{T}^S = T^S$.

If S is a segment of T^S and $f^{-1}(S)$ is not a subset of T^S , we can take a stable f -extension $\hat{T}$ with $\hat{T}^S = T^S \cup f^{-1}(S)$. We say this extension is formed by taking a backward iterate of S . In a similar way, we can form extensions by taking backward iterates of arcs of T^S , or forward iterates of arcs of T^U . For these extensions, the endpoints of any curve of $\hat{T}$ lie in the set $X = \bigcup_{n=-\infty}^{\infty} f^n(T^V)$.

By taking infinitely many iterates of T^U and T^S , we obtain a tangle known as an *infinite extension*.

Definition 2.25 (Infinite extension) Let $(f; T)$ be a trellis map. Then the *infinite f-extension* of T is the tangle $W(f; T^P)$.

As long as no point of T^P is the end of a curve of T , the tangle $W(f; T^P)$ can be obtained by iterating all branches of T infinitely often under f .

The main difference between extensions and supertrellises is that $\hat{T}^P = T^P$ for an extension, but T^P may be a strict subset of $\hat{T}^P$ for a supertrellis. This difference makes the analysis of supertrellises slightly more complicated than that of extensions. In particular, a curve in $\hat{T}^U$ may contain more than one periodic point, in which case all periodic points lie in the same region of T , and we can isotope so the curve contains an attractor between any two periodic points. This property makes supertrellises particularly useful, since we can first construct some invariant curves, and then decide how many points of $\hat{T}^P$ they should contain.




Figure 7: Extensions of the Smale horseshoe trellis (a) is a minimal extension, (b) is not.

Of particular interest are *minimal* iterates, extensions and supertrellises, which introduce as few extra intersections as possible given the endpoints of the branches. The trellis shown in Figure 7(a) is a minimal extension of the Smale horseshoe trellis, but the trellis shown in Figure 7(b) is not, since the intersection points x_0 and x_1 can be removed.

2.5 Reducibility

The concept of reducibility for trellis mapping classes is similar to that for mapping classes of surface diffeomorphisms, but is complicated by the fact that it may not be possible to decompose a system into a number of invariant subsets of negative Euler characteristic, but may be possible to decompose the system into an attractor-repeller pair. This gives two slightly different notions of a reductions, namely an *invariant curve reduction* and an *attractor-repeller reduction*.

Definition 2.26 (Invariant curve reduction) A trellis mapping class $([f]; T)$ has an *invariant curve reduction* if there is a diffeomorphism $\tilde{f} \in ([f]; T)$ and a one-manifold C which is disjoint from T , invariant under $\tilde{f}$, and each component of the complement of C either contains a subtrellis of T , or has negative Euler characteristic. If the complement of C is disconnected, we have a *separating reduction*, otherwise we have a *non-separating reduction*.

Definition 2.27 (Attractor-repeller reduction) A trellis mapping class $([f]; T)$ has an *attractor-repeller reduction* if there is a diffeomorphism $\tilde{f} \in ([f]; T)$ and a one-manifold C such that C divides M into subsets A and B such that $\text{cl}(\tilde{f}(A)) \subset \text{int}(A)$, and each component of the complement of C is either contains a point of T^P , or has negative Euler characteristic.

In both cases, the set C is called a set of *reducing curves*. It is clear that reducibility is a property of the trellis type, and not just the trellis mapping class.

The essence of both these definitions is that the manifold simplifies into two subsets which we can treat independently. However, the invariant curve reduction allows for curves do not divide the manifold, but instead decrease its genus. In both cases, if we find a reduction, we cut along the reducing curves, to obtain a simpler problem, or a number of simpler problems, which we deal with independently. After cutting along the reducing curves of an invariant curve reduction, we immediately obtain a diffeomorphism. After cutting along the reducing curves of an attractor-repeller curve reduction, we need to glue on an annulus to each curve. After glueing, we extend the diffeomorphism $\tilde{f}$ to the annulus in such a way that the remaining boundary is an attractor/repeller. For most of this paper, we only consider irreducible trellis mapping classes. We consider the problem of finding reductions in Section 5.




Figure 8: An trellis with an attractor-repeller reduction.

In Figure 8 we show a trellis type with an attractor-repeller reduction. A sub-basin A of an attractor is shaded. Notice that the unstable set $T^U(p_1)$ lies in A , which means it is possible to find a diffeomorphism $\tilde{f}$ for which $W^U(\tilde{f}; p_1) \cap W^S(\tilde{f}; p_0) = \emptyset$. The closure of the unstable manifold of p_1 is then a non-trivial chaotic attractor, though it may have non-trivial sub-attractors.

We now examine the hypothesis of irreducibility in more detail. Recall that the main effect of reducibility is that the dynamics of a reducible trellis mapping class can be separated into simpler pieces. We now show that the converse is also true; the dynam-

ics forced by an irreducible trellis essentially lies in one minimal component. To prove this, we need to carefully consider the intersections of branches of the trellis. We write $W^U(p_u, b_u) \times W^S(p_s, b_s)$ if the branches $W^U(p_u, b_u)$ and $W^S(p_s, b_s)$ intersect transversely.

For the case of trellises, we consider a relationship based on *heteroclinic chains*.

Definition 2.28 (Heteroclinic chains) Let $([f]; T)$ be an transverse trellis mapping class, and $p_u, p_s \in T^P$. We say there is a *heteroclinic chain* from p_u to p_s of length n if there are points $p_u = p_0, p_1, \dots, p_n = p_s$ in T^P and an integer m such that for every $\hat{f} \in ([f]; T)$ we have $T^U(p_i) \cap \hat{f}^{-m}(T^S)(p_{i+1}) \neq \emptyset$ for $i = 0, 1, \dots, n-1$. We write $p_u \succ p_s$ if there is a heteroclinic chain from p_u to p_s . A subset P of T^P is *chain transitive* if $p_1 \succ p_2$ for all $p_1, p_2 \in P$.

Clearly, $\succ$ is a transitive relation, though it need not be reflexive or symmetric.

Note that although $p_u \succ p_s$ implies $W^U(f; p_u)$ intersects $W^S(f; p_s)$ for any f in the trellis mapping class, it is not necessarily true that there exists an n such that $T^U(p_u)$ intersects $f^{-n}(T^S(p_s))$ for any f in the trellis mapping class. In other words, the intersections may lie in arbitrarily high iterates. For the next results, we therefore consider tangles rather than trellises.

Lemma 2.29 *Let $([f]; T)$ be a well-formed trellis mapping class, and $p_u = p_0, p_1, \dots, p_n = p_s$ be a heteroclinic chain. Let W be the infinite f -extension of T . Then $W^U(p_u) \times W^S(p_s)$.*

PROOF: By the Lambda lemma, if $W^U(p_{i-1}, b_{i-1}) \times W^S(p_i)$ and $W^U(p_i) \times W^S(p_{i+1}, b_{i+1})$, then $W^U(p_{i-1}, b_{i-1})$ transversely intersects $W^S(p_{i+1}, b_{i+1})$. Induction on the length of the heteroclinic chain completes the proof. $\square$

We now use the above observation to show that the chain transitive components of T are permuted.

Lemma 2.30 *Let $([f]; T)$ be a trellis mapping class. Suppose P is a chain transitive subset of T^P . Then there exists least $N \in \mathbb{N}$ such that $f^N(P) = P$. Further, if $0 < i < N$, then $p \not\succ f^i(p) \not\succ p$ for any i .*

PROOF: Suppose $p_u \succ p_s$. Let $\tilde{f}$ be a diffeomorphism such that $\tilde{f}^n(T^U(p_u)) \times T^S(p_s)$ and $(\tilde{f}^{n+1}(T^U), T^S)$ is a minimal extension of T . Then $\tilde{f}^{n+1}(T^U)(p_u) \times \tilde{f}(T^S(p_s))$, so $\tilde{f}^n(T^U(f(p_u))) \times T^S(f(p_s))$, so $f(p_u) \succ f(p_s)$. Similarly, we can see that $f^{-1}(p_u) \succ f^{-1}(p_s)$, and hence $f^n(p_u) \succ f^n(p_s)$ for any $n \in \mathbb{Z}$.

Since the points of P are periodic and reflexive under $\succ$, there exists least $N \in \mathbb{N}$ such that $p \succ f^N(p)$ for some $p \in P$. Then $f^{iN}(p) \succ f^{(i+1)N}(p)$, for any i , and since p is periodic, we deduce $p \succ f^{iN}(p) \succ p$ for any i .

Now suppose $q \in P$, so $p \succ q \succ p$. Then for any i , we have $p \succ f^{iN}(p) \succ f^{ni}(q) \succ f^{iN}(p) \succ p$, so $f^{iN}(q) \in P$. Conversely, we can show that if $f^{iN}(q) \in P$, then $q \in P$. Therefore, $P = f^N(P)$. $\square$

Note that the construction implies that N must divide the period of any point of P .

The relation $\succ$ induces a relation on the chain transitive components which is antisymmetric and transitive, but is not necessarily reflexive. If P is a reflexive component, then P contains a homoclinic orbit, so induces some chaotic behaviour. A chain transitive component P which is not reflexive must consist of a single periodic point p with $W^U(p) \cap W^S(p) = \emptyset$, so there are no homoclinic orbits to p .

If P_R and P_A are chain transitive components and $P_R \succ P_A \not\succ P_R$, then we hope to find an attractor-repellor reduction with P_A in the attractor and P_R in the repeller. If $P = f(P)$, then the tangle of the orbits of P is connected, but if $P = f^N(P)$ for some least $N > 1$, then the tangle of the orbits of P may have several components, which may form the basis for some invariant curve reduction. If P is not reflexive, then $W^U(P)$ may not intersect $W^S(P)$, and P is a non-chaotic saddle. The following theorem makes this interpretation precise.

Theorem 2.31 (Intersections of irreducible tangles) *Let $([f]; T)$ be a well-formed irreducible transverse trellis mapping class, and let W be the infinite f -extension of T . Then if $T^U(p_u, b_u)$ and $T^S(p_s, b_s)$ are closed branches of T , the extensions $W^U(p_u, b_u)$ and $W^S(p_s, b_s)$ intersect transversely.*

PROOF: The relationship $\succ$ clearly defines a partial order on the chain transitive sets. Choose a set P which is minimal in this partial order, and let $P_A = \bigcup_{n=1}^{\infty} f^n(P)$.

Let $([f_0]; T_0) = ([f]; T)$, and for $i = 1, \dots, \infty$, let $([f_i]; T_i)$ be a minimal extension of $([f_{i-1}]; T_{i-1})$ with $T_i = (f_i(T_{i-1}^U), T_{i-1}^S)$. Let K_i be a set consisting of $T_i^U(P_A)$, and all regions of T_i which are topological discs or annuli and all of whose stable boundary segments lie in $T^S(P_A)$. Since $T_i^U(P_A) \cap T_i^S(P_R) = \emptyset$, $K_i \subset K_{i+1}$. Let A_i be a closed neighbourhood of K_i which deformation retracts onto K_i , and such that $A_i \subset A_{i+1}$. Now, $\chi(K_i \setminus T^U) \geq \chi(K_{i+1} \setminus T^U) \geq \chi(M)$, so the Euler characteristics are a decreasing bounded sequence of integers, so have a limit, which occurs for some A_n . Then A_{n+1} deformation retracts onto K_n . Further, $f(K_n) \subset K_{n+1} \subset A_{n+1}$, so by taking A_n sufficiently small, we can ensure $f(A_n) \subset A_{n+1}$. Choose an isotopy h_t such that $h_1(A_{n+1}) \subset A_n$ and h_t is fixed on K , and let $\tilde{f}_t = h_t \circ f_n$. Then $\tilde{f}_1(A_n) = h_1(f_n(A_n)) \subset h_1(A_{n+1}) \subset A_n$, so $\tilde{f}_1 \in ([f]; T)$ and maps A_n into itself.

If $P_A \neq P$, then this gives an attractor-repellor reduction, a contradiction. If $P_A = P$ and $\chi(A_n) \neq \chi(M)$, we can isotope f so that the boundary of A_n maps to itself, giving an invariant curve reduction, also a contradiction. Therefore $P_A = P$ and there is no reduction. Since $T^U(p_u, b_u)$ and $T^S(p_s, b_s)$ are closed branches, there are periodic points $\tilde{p}_u$ and $\tilde{p}_s$ such that $T^U(p_u, b_u) \times T^S(\tilde{p}_u)$ and $T^U(\tilde{p}_s) \times T^S(p_s, b_s)$. Then $T^U(\tilde{p}_u) \times T^S(\tilde{p}_s)$, and so $T^U(p_u, b_u) \times T^S(p_s, b_s)$ as required. $\square$

In other words, if $([f]; T)$ is irreducible, then for there exists n such that the n th image of every unstable branch under f intersects every stable branch. This is, essentially, the converse to the trivial observation that if $([f]; T)$ has an attractor-repellor reduction, then there is a diffeomorphism $\tilde{f}$ in $([f]; T)$ and branches of $W(\tilde{f}; T^P)$ which do not intersect.

2.6 Cutting

Given a trellis, the first step in obtaining rigorous information on the dynamics is to *cut* along the unstable manifold to obtain a new space $\mathcal{C}T$, a topological pair which we denote $(\mathcal{C}^U T, T^S)$. A formal description of cutting is given in [Col99], so we shall not give one here as it is intuitively obvious.




Figure 9: Cutting along the unstable curve.

Cutting along the unstable curve of the Smale horseshoe trellis gives the set shown in Figure 9. The ends of the unstable curves gives *cusps* in the new surface. If T is a proper trellis, the stable segments lift to cross-cuts in the new surface $\mathcal{C}^U T$. End intersections of T^S also give isolated points in $\mathcal{C}^U T$ which are discarded. The topology of the lift of T^S to $\mathcal{C}^U T$ is sufficiently similar to that of T^S that we also denote this lift by T^S .

If f is a trellis map, $f(T^U)$ covers T^U , so we have a well-defined function $\mathcal{C}f$ on $\mathcal{C}^U T$. This function is not a diffeomorphism since a point q of T^U which maps to a point $f(q) \notin T^U$ will lift to two points with the same image. However, $\mathcal{C}f$ is surjective and at most two-to-one; in particular, f has the same topological entropy. Since $\mathcal{C}f$ maps T^S into itself, so is a map of the topological pair.

Notice that the pair $\mathcal{C}T = (\mathcal{C}^U T, T^S)$ contains the pair $(M \setminus T^U, T^S \setminus T^U)$ as an open subset which is invariant under $\mathcal{C}f$. Indeed, $\mathcal{C}^U T$ can be regarded as a natural compactification of $M \setminus T^U$. This compactness is important in the rigorous application of the Nielsen theory in Section 6. However, the homotopy properties of $(\mathcal{C}^U T, T^S)$ and $(M \setminus T^U, T^S \setminus T^U)$ are essentially the same.

2.7 Curves

Our main tool for studying the geometry, topology and dynamics associated with trellis maps will be to consider curves embedded in the cut surface $(\mathcal{C}^U T, T^S)$. As this surface is a topological pair, our curves will be maps in this category, so a curve α in T is a mapping $\alpha : (I, J) \longrightarrow (\mathcal{C}^U T, T^S)$, where I is the unit interval $[0, 1]$, and J is a closed subset of I . The *path* of such a curve α is the set $\alpha(I)$.

Since the topological pair $\mathcal{CT}$ is obvious from the trellis, we will usually draw curves in $\mathcal{C}^U T$ as curves embedded in the original surface M , and, wherever possible, ensure these curves are disjoint from T^U . However, as long as a curve in M does not cross T^U , it lifts to a curve in $\mathcal{C}^U T$. This simplifies many of the diagrams, and hence clarifies the geometry of the situation.

For the most part, we are only interested in curves up to homotopy or isotopy, and we always take homotopies and isotopies of curves through maps of pairs. For the Nielsen theory and shadowing results of Section 6, we will always keep the endpoints fixed during the homotopy. For most other purposes, we only consider curves in $\mathcal{C}^U T$ whose endpoints lie in T^S and for which the set J consists of finitely many points including 0 and 1. In this case, we only consider homotopies of curves which restrict to isotopies on J , or in other words, we ensure that α_t is one-to-one on J . However, we do allow the endpoints to move, though they are of course restricted to a segment of T^S .

If $\alpha : (I, J) \longrightarrow (X, Y)$ is a curve and J contains $\{0, 1\}$ we say α has endpoints in Y ; if J equals $\{0, 1\}$ we say α has endpoints only in Y .

Reparameterising a curve does not change its path, but may change the set J which maps into T^S . This means that different parameterisations of the same path may not even be comparable under homotopy. However, we consider different parameterisations of the same curve as equivalent.

Definition 2.32 (Equivalence of curves) Curves $\alpha_1 : (I_1, J_1) \longrightarrow (X, Y)$ and $\alpha_2 : (I_2, J_2) \longrightarrow (X, Y)$ are *equivalent* if there is a homeomorphism $h : (I_1, J_1) \longrightarrow (I_2, J_2)$ with $h(J_1) = J_2$ such that $\alpha_1 \sim \alpha_2 \circ h$ as curves $(I_1, J_1) \longrightarrow (X, Y)$. (The homotopy may be taken relative to endpoints, as appropriate.)

To study a trellis mapping class $([f]; T)$, we consider the iterates of an exact curve $\alpha : (I, J) \longrightarrow (\mathcal{C}^U T, T^S)$ with endpoints in T^S under models of $([f]; T)$. We are especially interested in iterates with minimal number of intersections with T^S .

Definition 2.33 (Minimal iterate) Let $([f]; T)$ be a trellis mapping class, and $\alpha : (I, J) \longrightarrow \mathcal{CT}$ be a simple curve with endpoints in T^S and transverse intersections with T^S . Then a *minimal iterate* of α is a curve β which is homotopic to $\mathcal{C}f \circ \alpha$ relative to J and which minimises the number of intersections with T^S . We let $J' = \beta^{-1}(T^S)$, and consider β as a curve $(I, J') \longrightarrow \mathcal{CT}$.

Theorem A.11 shows that the curve β is only well-defined up to equivalence. We therefore obtain a well-defined map $f_{\min}$ on equivalence classes of curves given by $f_{\min}[\alpha] = [\beta]$. An example of the first two minimal iterates of a curve α_0 is given in Figure 10. Note that by $f_{\min}^n[\alpha]$ we mean $(f_{\min})^n[\alpha]$, and not $(f^n)_{\min}[\alpha]$, which may be different. By $f_{\min}(\alpha)$, we mean some curve in the homotopy class $f_{\min}[\alpha]$. Minimal iterates are closely related to minimal extensions of trellises.

In Figure 11 we show a trellis type which is not well-formed, and a curve $\alpha_0 : (I, J) \longrightarrow (\mathcal{C}^U T, T^S)$ with endpoints in T^S . The minimal iterate of α_0 has self-intersections; note that by our notion of isotopy for such curves, points in $\alpha_t(J)$ cannot cross each other. However, this is impossible for a well-formed trellis mapping class.




Figure 10: Curve α_0 and minimal iterates $\alpha_1 \in f_{\min}[\alpha_0]$ and $\alpha_2 \in f_{\min}[\alpha_1]$.




Figure 11: Curve α_0 and minimal iterate α_1 which self-intersects.

Lemma 2.34 *Suppose $([f]; T)$ is a well-formed transverse trellis mapping class, and $\alpha : (I, J) \longrightarrow (\mathcal{C}^U T, T^S)$ is a simple exact curve in $\mathcal{C}^U T$ with endpoints in T^S and transverse intersections. Then there is a minimal iterate $f_{\min}[\alpha]$ which is a simple curve.*

PROOF: Let K be any disc in $\mathcal{C}^U T$ such that the boundary of K consists of an arc of β with endpoints x_0 and x_1 , and a subinterval of a segment of T^S . If neither x_0 or x_1 lie in $f(S)$, then there is a curve homotopic to β with fewer intersections with T^S than β obtained by removing the intersections x_0 and x_1 .

Suppose there is a point $q \in \alpha(J)$ such that $f(q) \in T^S(x_0, x_1)$, as shown in Figure 12, and let S be the component of T^S containing q . Then q lies in a component S of T^S . Then




Figure 12: If neither x_0 or x_1 lie in $f(T^S)$, then T is not well-formed.

$$f^{-1}(T^U) \cap S = f^{-1}(T^U \cap f(S) \subset f^{-1}(T^U \cap T^S(x_0, x_1)) = \emptyset,$$

which contradicts the fact that T is well-formed. Therefore, if there are any points of β in $T^S(x_0, x_1)$, these also do not lie in $\beta(J)$, so can be removed without introducing any self-intersections. $\square$

This result is a *homotopy* result, unlike the isotopy results of Section 7.

Sometimes we want to relate a curve $\hat{\alpha} : (\hat{I}, \hat{J}) \longrightarrow (\mathcal{C}^U T, T^S)$ to a curve $\alpha : (I, J) \longrightarrow (\mathcal{C}^U T, T^S)$. If $(\hat{I}, \hat{J}) \neq (I, J)$, the curves cannot be homotopic, but if $(I, J) \subset (\hat{I}, \hat{J})$ then the curve $\hat{\alpha}$ may be in some sense more complicated than α , as defined below.

Definition 2.35 (Tightening curves) Let $\hat{\alpha} : (\hat{I}, \hat{J}) \longrightarrow (X, Y)$ and $\alpha : (I, J) \longrightarrow (X, Y)$ be exact curves such that $(I, J) \subset (\hat{I}, \hat{J})$. We say $\hat{\alpha}$ *tightens to* α if the curve $\alpha_0 : (I, J) \longrightarrow (X, Y)$ given by $\alpha_0(t) = \hat{\alpha}(t)$ is homotopic to $\alpha_1 = \alpha$.

It is clear that the tightening relation is a partial order on homotopy classes of curves.

3 Biasymptotic Orbits and Minimal Trellises

We use the term *biasymptotic orbit* to refer to an orbit which is either homoclinic or heteroclinic to the set of periodic points. In this section, we consider how to extend a set of biasymptotic orbits to a trellis, and how to extend a trellis to a larger trellis. We shall usually consider extensions with a given *Birkhoff signature*. As previously remarked, this does not guarantee a unique trellis type. Instead, we define what it means for a trellis to be *minimal* given a biasymptotic mapping class. By carefully isotoping a given trellis map, we can construct a minimal trellis, and we then show that the type of a minimal trellis is almost unique given its end points, the only ambiguities arising from the orientation (transverse or tangent) of the intersections.

We also consider minimal iterates and minimal extensions of a given trellis mapping class. A minimal extension is forced by the biasymptotic orbits of the intersections of the original trellis. Unlike a minimal trellis for a biasymptotic mapping class, the orientations of the intersections are uniquely determined. We also define minimal supertrellises, for which we need to consider changes to the periodic point set.

3.1 Biasymptotic orbits

Just as we can consider isotopy classes of surface diffeomorphisms relative to periodic orbits and trellises, we can also consider isotopy classes relative to homoclinic and heteroclinic orbits to periodic saddle orbits.

Definition 3.1 (Biasymptotic mapping class) Let f be a surface diffeomorphism, and X a closed, invariant set consisting of a set of periodic saddle orbits X^P and a set X^V of biasymptotic orbits to X^P . The isotopy class of f relative to X is called the

biasymptotic mapping class $([f]; X)$. The conjugacy class of a biasymptotic mapping class is a *biasymptotic type*.

Methods for computing the dynamics forced by a biasymptotic mapping class have been given by Handel [Han99] and Hulme [Hul00]. Here we show how to relate a biasymptotic type to a trellis type class, and hence derive a new method for computing the dynamics.

Since X^P is a finite set of saddle orbits, we can compute the stable and unstable manifolds $W^{U/S}(f; X^P)$. Since the points of X^V are biasymptotic to X^P , they are all intersection points of W^U and W^S . However, W^U and W^S may have extra intersections which are not in X , and indeed, typically have infinitely many such intersections. Different representatives of a biasymptotic mapping class will have different tangles.

Definition 3.2 (Compatible tangle/trellis) Let $([f]; X)$ be a biasymptotic mapping class. Then if $\widehat{f}$ is any diffeomorphism in $([f]; X)$, the tangle $\widehat{W} = W(\widehat{f}; X^P)$ is a *compatible tangle* for $([f]; X)$. A trellis map $(\widehat{f}; \widehat{T})$ is *compatible* with $([f]; X)$ if $\widehat{f} \in ([f]; X)$ and $\widehat{T}$ is a subtrellis of the compatible tangle $\widehat{W}$ with $X^P = T^P (= W^P)$.

There are many different trellis types compatible with a biasymptotic type $([f]; X)$. To restrict the possible trellises, we consider only trellises with connected branches, and specify the endpoints of each branch as a point of X . This gives rise to the notion of a *Birkhoff signature*.

Definition 3.3 (Birkhoff signature) A *Birkhoff signature* for a biasymptotic mapping class $([f]; X)$ is a pair $\mathcal{B} = (\mathcal{B}^U, \mathcal{B}^S)$ of subsets of X such that $\mathcal{B}^U$ contains one point of X from each branch of $W^U(f; X^P)$ and $\mathcal{B}^S$ contain one point of X from each branch of $W^S(f; X^P)$. If $\mathcal{B}_1$ and $\mathcal{B}_2$ are two signatures, we say $\mathcal{B}_1 \leq \mathcal{B}_2$ if the trellis T_1 for f compatible with $(f; X)$ and with endpoints in $\mathcal{B}_1$ is a subtrellis of the compatible trellis T_2 with endpoints in $\mathcal{B}_2$.

We can further restrict the compatible trellises by specifying whether the intersection of W^U and W^S is transverse $\times$ or tangential $\asymp$ at each point of X . Note that all points on the same orbit must have the same intersection type.

Figure 13 shows some trellises compatible with the biasymptotic mapping class $([f]; X)$, where

$$X = \{p\} \cup \{q_i\} \cup \{v_i\}$$

The orbits q_i and v_i shown in Figure 13(a) are orbits of the Smale horseshoe map homoclinic to the saddle point p . Both these orbits lie in the same branch of $W^U(p)$ and $W^S(p)$, so a signature consists of a pair of points from either orbit. The trellis in Figure 13(b) has signature $(\{q_0\}, \{v_2\})$ and no intersections other than p . The trellises in Figure 13(c) and Figure 13(d) have signature $(\{q_2\}, \{v_0\})$. However, the trellis in Figure 13(c) has a tangency at v_1 , whereas the trellis in Figure 13(d) has a transverse intersection with negative orientation. The trellis in Figure 13(e) has signature $(\{v_2\}, \{q_0\})$, but has a pair of transverse intersections which is not forced by X .

If we specify the orientation of a crossing, it is not necessarily true that every compatible trellis map lies in the same isotopy class relative to the oriented intersections. An example




Figure 13: (a) Orbits $\{q_i\}$ and $\{v_i\}$ of the Smale horseshoe map. (b-e) Trellises compatible with $\{q_i\}$ and $\{v_i\}$.

is shown in Figure 14. Both trellis types shown have a set X consisting of the fixed saddle point p and the orbit of a homoclinic point v , and both yield the same biasymptotic mapping class $(f; X)$. Further, the orientation of the crossing at v is the same in each case. However, there is no isotopy between the maps in the different isotopy classes relative to X such that the orientation of the intersection at v does not change.

Clearly, even given a Birkhoff signature, there are still many, indeed infinitely many, trellis types compatible with a biasymptotic type. However, some trellises have extra intersections which can be removed by an isotopy in the biasymptotic mapping class. A trellis with no extra intersections is called a *minimal compatible trellis*. However, it is easiest to give a *local* definition which characterises the properties of the trellis more precisely, and from which minimality of the number of intersections follows.

Definition 3.4 (Minimal trellis) Let $([f]; X)$ be a biasymptotic mapping class. Then a trellis map $(\tilde{f}; \tilde{T})$ is a *minimal compatible trellis map* if $\tilde{f} \in ([f]; X)$ and

1. The endpoints of $\tilde{T}^U$ and $\tilde{T}^S$ are points of X .
2. Every bigon of $\tilde{T}$ either contains a point of X in the interior of one of its boundary segments, or both vertices are points of X .
3. Every tangency of $\tilde{T}^U$ and $\tilde{T}^S$ is a point of X .

The trellis $\tilde{T}$ is called a *minimal trellis* for $([f]; X)$.

We can also extract the biasymptotic orbits from a trellis. Clearly, if $(f; T)$ is a trellis map, the set

$$X = \bigcup_{n=-\infty}^{\infty} f^n(T^V)$$




Figure 14: The trellis mapping types shown in (a) and (b) are not isotopic through maps preserving the intersection at v .

is a set of biasymptotic orbits, and T is a minimal compatible trellis with $([f]; X)$. However, there may be smaller sets of biasymptotic orbits for which T is a minimal compatible trellis. Such a set is called a set of *forcing orbits*.

Definition 3.5 (Forcing) Let $(f; T)$ be a proper trellis map. A set X is a set of *forcing orbits* for T if T is a minimal compatible trellis for $([f]; X)$. We say that T is *forced* by X .

The rest of this section is mostly devoted to proving the existence and (essential) uniqueness of minimal trellises.

3.2 Existence of minimal trellises

To find a minimal trellis $\tilde{T}$ for a biasymptotic mapping class $([f]; X)$ with a given Birkhoff signature $\mathcal{B}$, we start with any compatible trellis map $(f_0; T_0)$ and try to remove intersection points by isotopy of maps in $([f]; X)$. Such an isotopy is called a *pruning isotopy*.

Definition 3.6 (Pruning isotopy) Let $([f]; X)$ be a biasymptotic mapping class, and $\mathcal{B}$ be a Birkhoff signature for $([f]; X)$. Let f_t be an isotopy of maps in $([f]; X)$, and T_t be the trellis for f_t with signature $\mathcal{B}$. Then $(f_t; T_t)$ is a *pruning isotopy* if for every open set U of the surface M , and every $\tau \in [0, 1]$, there is a neighbourhood V of τ such that the number of intersections of T_t in U decreases as t increases in V ,

In other words, no extra local intersections are created in T_t as t increases through a pruning isotopy. An example of a pruning isotopy is shown in Figure 15.

Assuming we always have finitely many intersections, the trellis type only changes at a tangency of T_t^U and T_t^S . Passing through such a tangency allows us to remove a pair of intersection points. These intersection points must be the vertices of a special kind of bigon called an *inner bigon*.




Figure 15: A pruning isotopy starting with the Smale horseshoe trellis (a) and ending with a minimal trellis relative to the marked point (f).

Definition 3.7 (Inner bigon) A bigon B of a trellis T with connected branches is *inner* if there exist positive integers n_u and n_s such that both $f^{n_u}(B)$ and $f^{-n_s}(B)$ are bigons, but $f^{n_u+1}(B^U) \cap T^U = \emptyset$ (or is an end of T^U) and $f^{-(n_s+1)}(B^S) \cap T^S = \emptyset$, where B^U and B^S are respectively the open unstable and stable boundary segments of B . A *first inner bigon* is an inner bigon for which $n_s = 0$, and a *last inner bigon* is an inner bigon for which $n_u = 0$,

Essentially, an inner bigon has no bigons inside it or any iterate. In particular, the sets $f^n(B)$ are disjoint for $-n_s \leq n \leq n_u$. Clearly, every inner bigon is the image of some first inner bigon. Note that the conditions preclude an inner bigon with a vertex in T^P .




Figure 16: The Smale horseshoe trellis with bigons B_0 and B_1 .

In Figure 16 we show a trellis with two labelled bigons, B_0 and B_1 . The bigon B_1 is a first inner bigon, but B_0 is not an inner bigon since $f^{-1}(B_0)$ is a domain but not a region.

The following lemma shows that if we have a trellis map with a bigon for which neither vertex is a periodic point, there must be an inner bigon.

Lemma 3.8 *Suppose $(f; T)$ is a trellis map, where T is a proper trellis with connected branches, and let B be a bigon which does not contain a point of T^P . Then T has an inner bigon.*

PROOF: Let B be any bigon, and let n_s be the greatest integer such that $f^{-n_s}(B_0^S)$ is bounded by an interval in T^U and an interval in T^S . Then $f^{-n_s}(B)$ contains no stable curves in its interior, but may contain unstable curves. Since $f^{-n_s}(B)$ is a topological disc, it must either be a bigon, or contain a bigon B as a subset. Then there exists a least integer n_u such that $f^{n_u+1}(B^U) \cap T^U = \emptyset$. Since B is a bigon, $n_u < n_s$, and therefore $f^{n_u}(B) \cap T^S \subset f^{n_u-n_s}(B) \cap T^S \subset f^{n_u-n_s}(B^S) \subset f^{n_u}(B^S)$, so $f^{n_u}(B)$ is also a bigon. Hence B is an inner bigon. $\square$

Henceforth we will only consider pruning isotopies which preserve the unstable set T^U , but allow us to change the stable set T^S . Unfortunately, changes made to one piece of stable manifold may affect other pieces. Therefore, we only consider of the form $f_t = f \circ h_t^{-1}$, where h_t is supported on a set K whose iterates are controlled as follows.

Definition 3.9 (Simple pruning disc) Let $([f]; T)$ be a trellis mapping class. An open disc K is a *simple pruning disc* if

1. $T^U \cap f(K) = T^S \cap f^{-N}(K) = \emptyset$,
2. $T^U \cap f^{-N}(K) = f^{-N}(T^U \cap K)$, and
3. the sets $K_n = f^{-n}(K)$ are disjoint for $0 \leq n < N$.

The following lemma shows how precomposing with the inverse of a diffeomorphism h supported in a simple pruning disc changes the stable and unstable manifolds.

Lemma 3.10 *Let $([f]; T)$ be a trellis mapping class with signature $\mathcal{B}$, and K be an open subset of M such that Let h be a diffeomorphism supported on K such that h is the identity on orbits of $\mathcal{B}$. Let $\tilde{f} = f \circ h^{-1}$, and $\tilde{T}$ be the trellis of $\tilde{f}$ with signature $\mathcal{B}$. Then $\tilde{T}^U = T^U$, $\tilde{T}^S = T^S$ outside of $\bigcup_{n=0}^{N-1} f^{-n}(K)$, and $\tilde{T}^S = h(T^S)$ in K .*

PROOF: Since $f^{-1}(T^U) \cap K = \emptyset$, if $x \in T^U$, then $f_t^{-1}(x) = h_t^{-1}(f^{-1}(x)) = f^{-1}(x)$, so $x \in \tilde{T}^U$. Hence $\tilde{T}^U = T^U$.

$f^N(T^S) \cap K = f^N(T^S \cap f^{-N}(K)) = \emptyset$. Therefore, if $x \in T^S$ and $f^n(x) \notin K$ for $0 \leq n < N$, then x never enters K , so $f^m(x) = \tilde{f}^m(x)$ for all m . Hence $x \in \tilde{T}^S$.

If $1 \leq n < N$, then

$$f^n(T^S \cap K) \cap K \subset f^n(K) \cap K = \emptyset,$$

and if $N \leq n$, then

$$f^n(T^S \cap K) \cap K \subset f^n(T^S) \cap K \subset f^N(T^S) \cap K = \emptyset.$$

Therefore, if $x \in h(T^S) \cap K$, we have $\tilde{f}(x) = f(h^{-1}(x)) \in f(T^S \cap K)$ since $h(T^S) \cap K = h(T^S) \cap K$. So $\tilde{f}^n(x) = f^n(h^{-1}(x))$ for any $n > 0$ since $f^n(h^{-1}(x)) \notin K$. Hence $x \in \tilde{T}^S$. $\square$

Notice that since K is a disc, the diffeomorphism h is isotopic to the identity, so $\tilde{f}$ is isotopic to f .

Neighbourhoods of inner bigons and points of T^V give suitable simple pruning discs.

Lemma 3.11 *Let $([f]; T)$ be a trellis map compatible with a biasymptotic mapping class $([f]; X)$. If B is a last inner bigon such that no point of B lies in X , then there is a simple pruning disc $K \supset B$. Similarly, if x is a point of T^V such that $f(x) \notin T^U$, there is a simple pruning disc $K \ni x$.*

The proof is straightforward from the definitions of inner bigon and simple pruning disc.

We can now give the main result of this section

Theorem 3.12 (Existence of minimal trellis) *Suppose $([f]; X)$ is a biasymptotic mapping class and $\mathcal{B} = (\mathcal{B}^U, \mathcal{B}^S)$ is a signature in X . Then there is a minimal trellis map $(f; \tilde{T})$ compatible with $([f]; X)$ with signature $\mathcal{B}$.*

PROOF: Suppose $(f; T)$ is not minimal. Then either T has a tangency v which is not in X , or by Lemma 3.8 we can find a last inner bigon B such that at least one vertex of B is not a point of X .




Figure 17: Isotopies removing intersections in K : (a) removing intersections from a bigon, (b) removing a tangency, (c) making a tangency at a point of X and (d) moving a transverse intersection to a point of X .

If v is a tangency point which is not in X , then there exists a greatest integer n_u such the $f^{n_u}(v) \in T^U$. By Lemma 3.11 there is a simple pruning disc K containing $f^{n_u}(v)$ and no other intersections of T . Take h such that $h(T^S \cap K)$ does not intersect T^U

and $\tilde{f} = f \circ h^{-1}$. Then by Lemma 3.10, $\tilde{T}$ has fewer intersection points than T . Then $f^{n_u}(B) \cap T^U = f^{-(n_s+1)}(B) \cap T^S = \emptyset$ since the end vertices of T are points of X . This case is shown in Figure Figure 17(a).

Similarly, suppose T has a last inner bigon B such that neither vertex of B is a point of X . Then by Lemma 3.11, there is a simple pruning disc K containing B , and no points of T^V other than the vertices of B , so we can isotopy f to $\tilde{f}$ with a trellis $\tilde{T}$ which has fewer intersection points than T . Then $f^{n_u}(B) \cap T^U = f^{-(n_s+1)}(B) \cap T^S = \emptyset$ since the end vertices of T are points of X . Since the sets $f^n(B)$ are disjoint for $-n_s \leq n \leq n_u$, we can find a contractible neighbourhood K of $f^{n_u}B$ satisfying the conditions of Lemma 3.10. Then, as before, we can find isotope f to obtain a trellis with fewer intersections. This case is shown in Figure Figure 17(b).

It remains to consider the cases where there is a last inner bigon B for which at least one vertex x is a point of X . Now, it may be the case that $f(x) \in T^U$ or $f^{-(n_s+1)}(x) \in T^S$, so Lemma 3.10 does not apply directly to B . In this case, we can first precompose with an isotopy supported in a neighbourhood of $f(x)$ to obtain an inner bigon whose vertices are not points of X , as shown in Figure 17(c) and Figure 17(d), and then remove this inner bigon as previously discussed. This again reduces the number of intersection points.

The result follows by induction on the number of vertices, since we can always obtain a trellis with finitely many vertices by perturbing f . Note that this construction does not change the unstable set T^U . $\square$

3.3 Uniqueness of minimal trellis

We now show that a minimal trellis compatible with a given biasymptotic mapping class with a given signature is essentially unique. Unfortunately, there is one problem; if there are two points x_0 and x_1 of X which form the intersections of an inner bigon, the orientation of these intersections is unspecified by the minimality conditions. Note that if there is a bigon with both vertices in X , we may have different local behaviour as shown in Figure 18.



Figure 18: Possibilities for bigons with both vertices in X .

As shown in Figure 19, there are minimal compatible trellises with transverse and tangential crossings at x_0 and x_1 .

We use the following preliminary theorem, which is derived from Theorem A.8 and Theorem A.11.




Figure 19: Different minimal compatible trellises.

Proposition 3.13 *Suppose $(f_0; T)$ and $(f_1; T)$ are minimal trellis maps compatible with a biasymptotic mapping class $([f]; X)$. Then $(f_0; T)$ and $(f_1; T)$ are in the same trellis mapping class.*

PROOF: Let $h = f_0^{-1} \circ f_1$. Then h is isotopic to id relative to X . Since T^U consists of cross-cuts to X , there is an isotopy $\tilde{h}_t$ such that $\tilde{h}_0 = id$, $\tilde{h}_1 = h$ and $\tilde{h}_t(T^U) = T^U$ for all t . If we now cut along T^U , we find that h is isotopic to the identity by the isotopy $\tilde{h}$, and T^S is a set of cross-cuts. Therefore, there is an isotopy h_t such that $h_0 = id$, $h_1 = h$, and for all values of t , $h_t(T^U) = T^U$ and $h_t(T^S) = T^S$. To show that $f_t = f_0 \circ h_t$ gives the required isotopy from f_0 to f_1 , we need only check that f_t preserves T^U and T^S . We have $f_t^{-1}(T^U) = h_t^{-1}(f_0^{-1}(T^U)) = f_0^{-1}(T^U)$ since $f_0^{-1}(T^U) \subset T^U$, and $f_t(T^S) = f_0(h_t(T^S)) = f_0(T^S)$, as required. $\square$

This result is important in its own right, since it shows that as long as the topology of the surface M is simple, we can specify a trellis mapping class by giving the trellis up to homeomorphism, and then only need to give one biasymptotic orbit on each branch. This greatly simplifies the representation of a trellis mapping class required for computations, and justifies our custom of specifying a trellis mapping class by drawing the trellis and part of a biasymptotic orbit.

We can now show that a minimal trellis compatible with a set of biasymptotic orbits is essentially unique.

Theorem 3.14 (Uniqueness of minimal trellis) *Let $([f]; X)$ be a biasymptotic mapping class and $\mathcal{B}$ a Birkhoff signature for X . Let $(f_0; T_0)$ and $(f_1; T_1)$ be trellis maps forced by $([f]; X)$, with end intersections at points of $\mathcal{B}$. Suppose that whenever there are points x_1 and x_2 of X such that $T^U[x_1, x_2] \cup T^S[x_1, x_2]$ bounds a bigon B in T_0 , and there are no other points of X on the boundary of B , that T_0 and T_1 have the same type of crossing (transverse or tangential) at x_1 and x_2 . Then $(f_0; T_0)$ and $(f_1; T_1)$ have the same trellis type.*

PROOF: Since the curves T^U and T^S are mutually homotopic, have the same types of intersection at X , and have minimal intersections with respect to X , by Theorem A.11, there is a homeomorphism h such that $h(T_0) = T_1$ and h is isotopic to the identity relative to X . Then $(h^{-1} \circ f_1 \circ h; h^{-1} \circ T_1) = (h^{-1} \circ f_1 \circ h; T_0)$, and f_0 and $h^{-1} \circ f_1 \circ h$ are isotopic relative to X . Then by Proposition 3.13, $(f_0; T_0)$ and $(h^{-1} \circ f_1 \circ h; T_0)$ are in the same

trellis mapping class, so $(f_0; T_0)$ and $(f_1; T_1)$ have the same type. $\square$

3.4 Minimal extensions

We have considered how to construct a from a set of biasymptotic orbits a trellis with a given signature. We can also extend trellises to longer trellises. Of particular interest are those which do not introduce any more intersections than necessary. Such trellises are called *minimal iterates* and *minimal extensions*. They are most simply characterised as being extensions which are minimal with respect to the orbits of the original trellis.

Definition 3.15 (Minimal iterate and minimal extension) Let $([f]; T)$ be a trellis mapping class. An iterate $(\tilde{f}; \tilde{T})$ of $([f]; T)$ is a *minimal iterate* of $([f]; T)$ if $\tilde{T}$ is forced by X , where the set X is given by $X = \bigcup_{n \in \mathbb{Z}} \tilde{f}^n(T^V)$, the union of the orbits of the intersections of T . An extension $(\tilde{f}; \tilde{T})$ of $([f]; T)$ is a *minimal extension* of $([f]; T)$ if it is a subtrellis of some minimal iterate.

The following result follows almost directly from the definition of a minimal extension.

Theorem 3.16 (Existence and uniqueness of minimal extensions) Let $([f]; T)$ be a trellis mapping class, $X = \bigcup_{n \in \mathbb{Z}} f^n(T^V)$, $\mathcal{B}$ the signature class of T in X , and $\tilde{\mathcal{B}}$ a signature class in X with $\tilde{\mathcal{B}} \geq \mathcal{B}$. Then there is a unique trellis type $[\tilde{f}; \tilde{T}]$ with signature $\tilde{\mathcal{B}}$ which is a minimal extension of $([f]; T)$. Further, if T is a transverse trellis, so is $\tilde{T}$.

The proof is essentially the same as that of Theorem 3.12 and Theorem 3.14. Note that any inner bigon of $\tilde{T}$ is an iterate of some inner bigon of T , so the intersection type is given by T .

The following trivial result gives a characterisation of certain minimal extensions.

Lemma 3.17 Let $(\tilde{f}; \tilde{T})$ be a minimal extension of $([f]; T)$. Then every bigon of $\tilde{T}$ contains a point of $\tilde{X} = \bigcup_{n=-\infty}^{\infty} f^n(T^V)$. Further, if $(\tilde{f}; \tilde{T})$ is any extension for which the end intersections in $\tilde{X}$, and every bigon of $\tilde{T}$ contains a point of $\tilde{X}$, then $(\tilde{f}; \tilde{T})$ is a minimal extension.

Note that it is possible to find extensions of $([f]; T)$ for which every bigon contains a point of $\tilde{X}$, but which is not a minimal extension. By Lemma 3.17, this may only occur if there is an end intersection not in $\tilde{X}$.

The definition of a minimal supertrellis is complicated since we need to ensure that the extra periodic points correspond to orbits of the original trellis mapping class.

Definition 3.18 Let $([f]; T)$ be a trellis mapping class. Then a supertrellis $(\tilde{f}; \tilde{T})$ is a *minimal supertrellis* if for every extension of $(\tilde{f}; \tilde{T})$ of $([f]; \tilde{T})$, every bigon of $\tilde{T}$ contains a point of $\tilde{X} = \bigcup_{n=-\infty}^{\infty} \tilde{f}^n(T^V)$.

In particular, this means that every point of $\tilde{T}^P \setminus T^P$ shadows an essential periodic orbit of $([f]; T)$.

Since minimal extensions and minimal supertrellises have the same forcing orbits as the original trellis, we expect the dynamics to be the same as that forced by the original trellis, or indeed, the dynamics forced by the biasymptotic orbits themselves. This is in fact the case, as we shall see later. Minimal extensions provide a way of getting more information about the dynamics without increasing entropy.

4 Graphs Representatives

Any connected compact surface with nonempty boundary is homotopy-equivalent to a one-dimensional space. Further, such spaces, and maps on them, are very easy to describe combinatorially. We can use these properties to provide a framework for representing surface diffeomorphisms and computing their dynamical properties. In classical Nielsen-Thurston theory, surface homeomorphisms are represented on one-dimensional spaces with a differentiable structure called *train tracks*. When representing trellises, we also need to take into account the topology of the stable and unstable curves. By cutting along the unstable curves, we introduce new loops in the graph. The stable curves are represented by special edges called *control edges*, since these control the behaviour of the one-dimensional representative of a trellis map.

4.1 Combinatorics of graph maps

We first give some standard definitions concerning graphs embedded in compact surfaces.

Definition 4.1 (Graph) A *graph* G is an one-dimensional CW-complex. The *vertices* of G are the zero-dimensional cells, and the *edges* are the one-dimensional cells. The edges are oriented, and the reverse of the edge e is denoted $\bar{e}$, so $\bar{\bar{e}} = e$. The initial vertex of a oriented edge e will be denoted $\iota(e)$. The *valence* of a vertex v is the number of directed edges for which v is the initial vertex. A graph can be described up to homeomorphism by giving its vertices and the incident edges at each vertex.

The *Euler characteristic* χ of a graph is $|V(G)| - |E(G)|$ where $|V(G)|$ is the total number of vertices and $|E(G)|$ is the total number of edges. χ is an homotopy type invariant.

Definition 4.2 (Edge-path) An *edge-path* is a list $e_1 \dots e_n$ of oriented edges of G such that $\iota(\bar{e}_i) = \iota(e_{i+1})$ for $1 \leq i < n$. An *edge loop* is a cyclically-ordered list of edges. The *trivial* edge-path contains no edges and is denoted $\cdot$. An edge-path $e_1 \dots e_n$ *back-tracks* if $e_{i+1} = \bar{e}_i$ for some i , otherwise it is *tight*.

Definition 4.3 (Surface embedding) We will always consider a graph embedded in a surface by an embedding i . This induces a natural cyclic order $\triangleright$ on the oriented edges starting at each vertex. The graph can be described up to local ambient homeomorphism by giving the cyclic ordering of the incident edges at each vertex.

Definition 4.4 (Turn) A pair of edges (e_1, e_2) is a *turn* in G at vertex v if $v = \iota(e_1) = \iota(e_2)$ and $e_2 \triangleright e_1$, so e_2 immediately follows e_1 in the cyclic order at v .

Definition 4.5 (Peripheral loop) An edge-loop $\pi = \dots, p_1, p_2, \dots, p_n, \dots$ is *peripheral* in G if $(p_{i+1}, \bar{p}_i)$ is a turn in G for all i . A peripheral loop π is *simple* if π is a simple closed curve, or equivalently, if $\iota(p_i) \neq \iota(p_j)$ for $i \neq j$. Peripheral loops which are not simple may repeat vertices, or even edges (with opposite orientation). Peripheral loops define anti-clockwise curves around boundary components of the surface M .

Definition 4.6 (Frontier and link) If H is a subgraph of G , an edge-loop π is a *frontier loop* of H if π is peripheral in H but not in G . If H is a subgraph of G , then the *link* of H , denoted $\text{Lk}(H)$ is the set of edges e of G with $\iota(e) \in H$, but $e \notin H$.

Definition 4.7 (Graph maps) A *graph map* g is a self-map of G taking a vertex to a vertex, and edge e to an edge-path $e_1 \dots e_k$ such that $\iota(e_1) = g(\iota(e))$ for all directed edges e . If g is a graph map on G , where G is embedded in M by an embedding i , then we say g is *embeddable* if there are arbitrarily small perturbations of $i \circ g$ which are embeddings and preserve the cyclic order at any vertex. (A combinatorial definition of an embeddable graph map also exists.)

Definition 4.8 (Derivative map) The *derivative* map ∂g takes oriented edges to oriented edges or $\cdot$, with $\partial g(e_i) = e_j$ if $g(e_i) = e_j \dots$ and $\partial g(e_i) = \cdot$ if $g(e_i) = \cdot$.

Definition 4.9 (Peripheral subgraph) The *peripheral subgraph* P of g is a maximal invariant subset of G consisting of simple peripheral loops. Edges of P are called *peripheral edges*. If $g^n(e) \subset P$ for some n , then e is *pre-peripheral*. The set of pre-peripheral edges is denoted $\text{Pre-}P$, and contains P .

The dynamics of g can be split into *transitive components*.

Definition 4.10 (Transitive and irreducible component) Define a transitive relation on the edges E of G by $e_1 \geq e_2$ if there exists an n such that $g^n(e_1)$ contains the edge e_2 , and a stronger transitive relation by $e_1 \succ e_2$ if there exists n such that $g^m(e_1)$ contains the edge e_2 for all $m \geq n$. A *transitive component* of g is an equivalence class under $\geq$, and an *irreducible component* of g is an equivalence class under $\succeq$.

The quotient of these relation by the equivalence classes give a partial orders on the transitive/irreducible components. A transitive component of a graph map g is a union of irreducible components which are cyclically permuted by g .

Definition 4.11 (Transition matrix and growth rates) The *transition matrix* of g is the matrix $A = (a_{ij})$ where a_{ij} is the number of times the edge e_j appears in the image path of edge e_i . The largest eigenvalue of A is the *growth rate* λ of g , and the logarithm of the growth rate gives the topological entropy of g .

We can compute growth rates for any transitive component, and the growth rate for the entire graph map is the maximum of the growth rates on each component.

Example 4.12

Figure 20 shows a graph embedded in a disc with four punctures. The peripheral loops are p_1, p_2, p_3, p_4p_5 (surrounding the punctures) and $a\bar{p}_1\bar{a}\bar{p}_4c\bar{p}_3\bar{c}\bar{p}_5b\bar{p}_2\bar{b}$.




Figure 20: A graph embedded in a disc with four punctures.

There is a graph map g such that

$$\begin{aligned} g(p_1) &= p_2, \quad g(p_2) = p_3, \quad g(p_3) = p_1, \quad g(p_4) = p_5, \quad g(p_5) = p_4, \\ g(a) &= c\bar{p}_3\bar{c}\bar{p}_5b, \quad g(b) = c\bar{p}_3\bar{c}\bar{p}_5b\bar{p}_2\bar{b}\bar{p}_5c, \quad g(c) = a. \end{aligned}$$

The peripheral subgraph of g is $\{p_1, p_2, p_3, p_4, p_5\}$. The link of the peripheral subgraph consists of the oriented edges $\{a, \bar{a}, b, \bar{b}, c, \bar{c}\}$.

When relating graph maps to trellis types in Section 4.2 we need to consider topological pairs to capture the configuration of T^S .

Definition 4.13 (Control edges) Let (G, W) be a topological pair, where W is a finite set of points such that every edge of G contains at most one point of W . Then the edges of G containing a point of W are called *control edges*; other edges of G are *free edges*. The set of control edges of G is denoted $Z(G)$. The pair (G, W) is a *controlled graph*. A graph map $g : (G, W) \rightarrow (G, W)$ is a *controlled graph map* if for every control edge z_0 , the image of $g(z_0)$ is a control edge z_1 . We denote a controlled graph map g of (G, W) by $(g; G, W)$.

A vertex of G which is the endpoint of a control edge is called a *control vertex*. All other vertices are called *free vertices*, and edges which are not control edges are called *free edges*. Free edges which are neither peripheral nor pre-peripheral are called *expanding edges*.

A controlled graph is *proper* if it is connected and free vertex has valence at least 3. A proper controlled graph has at most $3|C(G)| - 3\chi$ edges, and $3|C(G)| - 2\chi$ vertices, where $|C(G)|$ is the number of control edges.

4.2 Compatible graph maps

Our main tool for computing and describing the dynamics forced by a trellis mapping class or trellis type is to relate the trellis map to a graph map. We do not relate a trellis

mapping class directly to a graph map, but instead use the map $\mathcal{C}f$ induced by cutting along the unstable curves. To relate this map to a map on a different space, we introduce a new concept of equivalence, that of *exact homotopy equivalence*, which is the natural extension of the concept of homotopy equivalence to maps of pairs.

We first need to define what it means for a map of pairs to be *exact*.

Definition 4.14 (Exact map of pairs) A map of topological pairs $f : (A, B) \longrightarrow (X, Y)$ is *exact* if $f^{-1}(Y) = B$. (Equivalently, f is exact if $f(B^C) \subset Y^C$.)

Note that this is equivalent to saying that f is a map of triples, $f : (A, B, B^C) \longrightarrow (X, Y, Y^C)$. However, we do not use this notation, as we treat exact maps as a special class of map in the category of topological pairs. In particular, we will often take homotopies between exact maps and non-exact maps in the category of topological pairs.

We can compare systems on different spaces via exact homotopy equivalence.

Definition 4.15 (Exact homotopy equivalence) Topological pairs (A, B) and (X, Y) are *exact homotopy equivalent* if there are exact maps $p : (A, B) \longrightarrow (X, Y)$ and $q : (X, Y) \longrightarrow (A, B)$ such that $q \circ p \sim id_A$ and $p \circ q \sim id_X$. p and q are called *exact homotopy equivalences*. Maps $f : (A, B) \longrightarrow (A, B)$ and $g : (X, Y) \longrightarrow (X, Y)$ are *exact homotopy equivalent* if there is an exact homotopy equivalence $p : (A, B) \longrightarrow (X, Y)$ such that $p \circ f = g \circ p$.

Note that the homotopies used in the definition of exact homotopy equivalence can be taken through any map of pairs and not just exact maps.

A graph map representing the topology of a trellis via exact homotopy equivalence is called *compatible* with the trellis.

Definition 4.16 (Compatible graph map) Let (G, W) be a controlled graph. The (G, W) is *compatible* with a transverse trellis T if (G, W) and $(\mathcal{C}^U T, T^S)$ are exact homotopy equivalent by an embedding $i : (G, W) \longrightarrow (\mathcal{C}^U T, T^S)$, and G crosses T^S transversely. A controlled graph map g of (G, W) is *compatible* with the trellis type $[f; T]$ if the embedding i is an exact homotopy equivalence between g and $\mathcal{C}f$, and g and $\mathcal{C}f$ have the same orientation at points of W .

Note that the inclusion i induced a bijection between the regions of $\mathcal{C}T$ (and hence of T) with the regions of (G, W) , and that all compatible graphs are exact homotopy equivalent. We restrict to transverse trellises since a trellis with tangencies may not have a compatible controlled graph. It is always possible to find a topological pair (G, H) which is exact homotopy equivalent to $(\mathcal{C}^U T, T^S)$ for which G is a graph, but it may not be possible to take H to be a finite set of points.

The controlled graph shown in Figure 21 is compatible with the trellis T . The control edges are shown as thick green lines.

There are many controlled graph maps compatible with a trellis type $[f; T]$. To use graph maps to describe the dynamics forced by $[f; T]$, we need to define a subclass of controlled




Figure 21: The controlled graph (G, W) is compatible with the trellis T .

graph maps which have minimal entropy in the exact homotopy class. These graph maps are called *efficient*, by analogy with Nielsen-Thurston theory. Of the efficient graph maps, we show in Section 4.5 that there is at most one which is *optimal*, giving a canonical *graph representative* for the trellis type.

Efficient and optimal graph maps can be defined in terms of their actions on the turns of G . Some of following notions are used in the definitions, and all are useful in describing the algorithms.

Definition 4.17 (Bad turns) Let g be a graph map of (G, W) , and (e_0, e_1) a turn of G . Then

1. If $\partial g(e_0) \neq \partial g(e_1)$, the turn is a *good* turn.
2. If $\partial g(e_0) = \partial g(e_1)$ and at least one of e_0 and e_1 is a control edge, the turn is a *controlled* turn. If both e_0 and e_1 are control edges, the turn is *fully controlled*, if only one is, the turn is *half controlled*.
3. If $\partial g(e_0) = \partial g(e_1)$ and both e_0 or e_1 are free edges, the turn is a *bad* turn.
4. A bad turn (e_0, e_1) is *inefficient* if in addition there is an edge e such that $g^n(e) = \dots \bar{e}_0 e_1 \dots$

In order to find efficient graph maps, we pass through intermediate graphs where there are no obvious ways of reducing the dynamics locally. Such maps are called *tight*.

Definition 4.18 (Tight graph map) Let g be a graph map of a controlled graph G . Then g is *vertex tight* if there is no free vertex v such that the derivative map ∂g takes the same value for all oriented edges e with $\iota(e) = v$, and g is *edge tight* if for every edge e , the edge path $g(e)$ is nontrivial and does not back-track. g is *tight* if it is both vertex tight and edge tight.

Note that ∂g is constant at a vertex v if there is an edge e such that $g(e_i) = e \dots$ for all edges e_i at v .

Definition 4.19 (Efficient and optimal graph maps) A controlled graph map g is *efficient* if there are no inefficient turns, and *optimal* if there are no bad turns, and every invariant forest contains a control edge.

A *graph representative* of $[f; T]$ is then an optimal controlled graph map compatible with $([f]; T)$.

Definition 4.20 (Graph representative) A controlled graph map $(g; G, W)$ is a *graph representative* of an transverse trellis type $[f; T]$ if g is an optimal graph map which is compatible with $([f]; T)$.

The main theorem concerning the representation of trellis types by graph maps is stated below.

Theorem 4.21 (Existence and uniqueness of graph representatives) *Let $[f; T]$ be a proper trellis type with no invariant curve reduction. Then $[f; T]$ has a unique graph representative $(g; G, W)$. Further, if $[f_0; T_0]$ and $[f_1; T_1]$ are trellis types, the graph representatives $(g_0; G_0, W_0)$ and $(g_1; G_1, W_1)$ are homeomorphic, then $[f_0; T_0] = [f_1; T_1]$.*

We prove the existence of a graph representative in Section 5 by giving an algorithm to compute it, and uniqueness in Section 4.5. Since the maps $\mathcal{C}f_0$ and $\mathcal{C}f_1$ for different trellis types are not exact homotopy equivalent, it is trivial that different trellis types have different graph representatives. This means that the graph representative provides a convenient way of specifying a trellis type.

If we are interested in the dynamics forced by the trellis mapping class, we can restrict attention to the subgraph of the graph representative $(g; G, W)$ containing the nontrivial dynamics. This subgraph is called the *essential graph representative*.

Definition 4.22 (Essential graph representative) Let $(g; G, W)$ be the graph representative of a trellis type $[f; T]$. Then the *essential graph representative* of $[f; T]$ is the controlled graph map $(g; \overline{G}, \overline{W})$, where $\overline{G} = \bigcap_{n=0}^{\infty} g^n(G)$, and $\overline{W} = \overline{G} \cap W$.

We can further simplify the representation by collapsing all control edges to points. This gives the *topological graph representative*. The topological graph representative is an invariant of the forcing orbits.

Definition 4.23 (Topological graph representative) Let $(g; G, W)$ be the graph representative of a trellis type $[f; T]$. Then the *topological graph representative* of $[f; T]$ is the topological conjugacy class of graph map obtained by collapsing all control edges of the essential graph representative to points.

Example 4.24

The graph representatives for the trellis type shown in Figure 21 are shown in Figure 22.

The control edges map

$$z_0, z_1, z_2 \mapsto z_0; \quad z_3 \mapsto z_1; \quad z_4 \mapsto z_2; \quad z_5 \mapsto z_3; \quad z_6, z_7, z_8 \mapsto z_4;$$

and the expanding edges map

$$a_0 \mapsto a_0 \bar{z}_8 z_6 b_0 \bar{b}_1; \quad b_0 \mapsto b_1; \quad b_1 \mapsto b_2; \quad b_2 \mapsto a_0 \bar{z}_8 z_6 b_0; \quad d \mapsto a_0 \bar{z}_8 z_6 b_0 \bar{b}_1.$$




Figure 22: Graph representatives for the trellis type shown in Figure 21. (a) is the graph representative, (b) the reduced graph representative and (c) the topological graph representative.

The edges z_1, z_3, z_5, z_7 and d do not lie in $g^2(G)$ so are not part of the essential graph representative shown in Figure 21(b). Collapsing the control edges gives the graph shown in Figure 21(c), with

$$a_0 \mapsto a_0 b_0 \bar{b}_1; \quad b_0 \mapsto b_1; \quad b_1 \mapsto b_2; \quad b_2 \mapsto a_0 b_0.$$

Joining the edges a_0 and b_0 to give a single edge e_0 , and letting $e_1 = b_1$ and $e_2 = b_2$ gives the topologically conjugate graph map

$$e_0 \mapsto e_0 \bar{e}_1 e_1; \quad e_1 \mapsto e_2; \quad e_2 \mapsto e_0.$$

The transition matrix for this graph map is

$$\begin{pmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}$$

and the growth rate is given by the largest root of the characteristic polynomial $\lambda^3 - \lambda^2 - 2 = 0$. Numerically, the growth rate is 1.696, giving topological entropy $h_{top}(g) \approx 0.528$.

4.3 Properties of graph representatives

We now give some elementary properties concerning relationship between a trellis and a compatible graph, and of the dynamics of a graph representative. Our main tool will be to look at curves in $\mathcal{C}T$ with endpoints in T^S which cross T^S transversely. These curves can be projected to (G, W) by the exact homotopy retract $r : (\mathcal{C}^U T, T^S) \longrightarrow (G, W)$. Since the inclusion $i : (G, W) \longrightarrow (\mathcal{C}^U T, T^S)$ is an exact homotopy inverse for r , there is a natural equivalence between exact homotopy classes of curves in $(\mathcal{C}^U T, T^S)$ and in (G, W) .

The ends of T^U give rise to *cusps* in (G, W) , as shown in Figure 23. In Figure 23(a), the control edges z_0 and z_1 surround an end of T^U . Then $g(z_0)$ and $g(z_1)$ must be the same, unless the intersection point q is a point of T^P . In Figure 23(b), the point q is an endpoint of T^S . In this case, the initial edge of $g(a_1)$ is $g(z_1)$, and the turn is only half controlled.

If z_0 and z_1 form a cusp, then unless z_0 and z_1 are periodic, in which case the cusp forms at the end of an open branch of T^U , we have $g(z_0) = g(z_1)$. The only other possibility for a




Figure 23: Cusps at the end of an unstable curve. (a) shows a cusp where the last intersection is a transverse crossing. (b) shows a cusp where the last intersection is an endpoint of T^S .




Figure 24: Turns surrounding a bigon.

turn where both edges are control edges is shown in Figure 24(a). Here, the control edges z_0 and z_2 are parallel to the unstable boundary B^U of a bigon B whose stable boundary crosses a control edge z_1 . If $f(B^U) \cap T^U = \emptyset$, then $g(z_0) = g(z_1) = g(z_2)$. Removing the stable boundary segments S_2 crossing z_2 as shown in Figure 24(b) does not significantly change the topology of the graph; in particular, the valence-3 vertex remains a control vertex. However, if we also remove the stable boundary segment S_0 crossing z_0 , this vertex is no longer a control vertex, so the resulting graph map is no longer optimal. Indeed, the graph representative for the resulting trellis may not be surface embeddable, and its entropy need not equal $h_{top}[f; T]$. This is a consequence of the failure of Lemma 2.34 for minimal iterates of curves.




Figure 25: A minimal iterate of a curve in a graph.

Now consider the graph shown in Figure 25. The curve α_0 from w_3 to w_2 maps to α_1 from w_2 to w_1 . We shall see that if the curve α_0 does not back-track except in control

edges, and g is an efficient graph map, then $g \circ \alpha_0$ is a minimal iterate. This will allow us to prove the existence of (bi)asymptotic orbits directly from the graph map. Also, by relating minimal extensions to the graph map using curves isotopic to T^U , we can easily find properties of minimal extensions.

For each point p of T^P , the boundary component obtained by cutting along $T^U(p)$ consists of two smooth pieces. These pieces are homotopic to a curve in the graph called an *unstable-parallel curve*.

Definition 4.25 (Unstable-parallel curve) A simple exact curve $\alpha : (I, J) \rightarrow (\mathcal{C}^U T, T^S)$ with endpoints in T^S is *unstable-parallel* if there is an exact isotopy $\alpha_t : (I, J) \rightarrow (\mathcal{C}^U T, T^S)$ such that $\alpha_0 = \alpha$ and α_1 lies in T^U .




Figure 26: Graph compatible with the Smale horseshoe trellis.

In Figure 26 we show a graph compatible with the Smale horseshoe trellis. Passing anticlockwise around the unstable $T^U(p)$, we have two smooth curves, one curve α with the same orientation as $T^U(p)$ and lying to its right, and one curve β with opposite orientation to its left. These curves can be projected onto the graph representative (G, W) giving edge-paths $\alpha = z_3 \bar{z}_3 z_1 c \bar{z}_5 z_5 \bar{c} \bar{z}_1$ and $\beta = z_0 a \bar{z}_6 z_4 \bar{b} \bar{z}_1 z_1 b \bar{z}_4 z_6 \bar{a} \bar{z}_0$. Note that each edge is traversed twice. This is always the case if we traverse all boundary components of $\mathcal{C}T$, including those which are boundary components of M .




Figure 27: Curve α_1 is unstable isotopic to T^U , and α_1 follows a minimal extension.

One important feature about unstable-parallel curves is that if $([\tilde{f}]; \tilde{T})$ is a minimal unstable extension of $([f]; T)$ and α is an unstable-parallel curve for T , then $\tilde{f}_{\min}(\alpha)$ is unstable-parallel curve for $\tilde{T}$. Therefore, we can use unstable parallel curves to construct minimal extensions, as shown in Figure 27. An unstable-parallel curve in a graph G compatible with a trellis T is a subpath of a peripheral loop.

4.4 Reducibility of graph representatives

Just as for trellis mapping classes, some graph maps are *reducible* and can be split into simpler pieces.

Definition 4.26 (Reducible graph map) A controlled graph map g of (G, W) is *reducible* if G has an invariant subgraph H such that H does not contain any control edges and either

1. H has negative Euler characteristic. We call this an *n-component separating* reduction, where n is the number of components of H .
2. H is a union of n simple closed curves but is not a subset of P . We call this an *n-curve non-separating* reduction.

g has an *attractor-repellor* reduction if there is an invariant subgraph H such that

3. H and H^C both contain control edges.

If a controlled graph map compatible with an trellis mapping class has a reduction, we cut out the invariant subgraph and consider the resulting pieces separately. The reduction algorithm is described in Section 5.4.

If $[f; T]$ has a separating invariant curve reduction which gives rise to an invariant sub-surface with no components of T , then there cannot be a graph representative, since we cannot move bad turns in the corresponding invariant subgraph to control edges. Therefore, we can only hope to find graph representatives for a trellis type without invariant curve reductions. The next result gives a condition for a trellis mapping class to have an attractor-repellor reduction.

Theorem 4.27 (Reductions of compatible graph maps) *Let g be a proper graph map compatible with a well-formed trellis type $[f; T]$. If g has a proper invariant subgraph H which is disjoint from W and either has negative Euler characteristic, or is a union of non-peripheral circles. Then $[f; T]$ has an invariant curve reduction. If g has any other proper invariant subgraph which either has negative Euler characteristic or is a union of non-peripheral circles, then g has an attractor-repellor decomposition.*

PROOF: If H is in invariant subgraph which is disjoint from W , then we can take the boundary curves of H to be a set of reducing curves. If H is any other invariant subgraph, let $\overline{H} = \bigcup_{n=0}^{\infty} g^{-n}(H)$ and let $\hat{H}$ be the set of components of H which either contain points of H or contain control edges mapping into H . Since $g(H) \subset H$, we must have $g(\hat{H}) \subset \hat{H}$,

and further, $g(\partial\widehat{H}) \subset \partial\widehat{H}$. Therefore $\partial\widehat{H}$ is contained in free edges. We introduce control edges at points of $\partial\widehat{H}$, which corresponds to introducing extra stable curves T_N^S curves in T . The region bounded by T^U and T_N^S is then an attractor, which we cut out to give one component of the reduction. The complement of this region is a repellor. We can find the dynamics of this repellor by removing all control edges in $\widehat{H}$, which corresponds to removing all stable curves in A . We cut along these curves to obtain the attractor. $\square$




Figure 28: Graph of a trellis with an attractor-repellor reduction.

Example 4.28

Figure 28 shows the graph of a trellis with an attractor-repellor reduction. The control edges z_i map

$$z_0, z_1, z_2 \mapsto z_0; \quad z_3 \mapsto z_1; \quad z_4 \mapsto z_2; \quad z_5 \mapsto z_3; \quad z_6, z_7, z_8 \mapsto z_4;$$

and the other control edges map

$$\begin{array}{ccccccc} C_{-7}, C_{-6} \mapsto C_3; & C_{-5}, C_{-4}, C_{-3} \mapsto C_2; & C_{-2} \mapsto C_1; & C_{-1}, C_0 \mapsto C_0; \\ C_0, C_1, C_2 \mapsto C_{-0}; & C_3 \mapsto C_{-1}; & C_4 \mapsto C_{-2}; & C_5 \mapsto C_{-3}; & C_6, C_7, C_8 \mapsto C_{-4}; & C_9 \mapsto C_{-5}; \\ & C_{10} \mapsto C_{-6}; & C_{11}, C_{12}, C_{13} \mapsto C_{-7}. \end{array}$$

The boundary points of the invariant set A are marked with crosses, and the control edges not in A are those labelled z_i , and the control edges in A are labelled c_i . The trellis with this graph representative is the one given in Figure 8.

We now show that the every edge eventually maps over a peripheral edge or a control edge, or g is reducible.

Lemma 4.29 *Let $(g; G, W)$ be the graph representative of an irreducible trellis type $[f; T]$. Then if e is any edge of g , there exists n such that $g^n(e)$ contains a control edge or a peripheral edge.*

PROOF: Let $H = \bigcup_{n=0}^{\infty} g^n(e)$, an invariant subgraph of G . Suppose H contains no control edges or peripheral edges. Then either H is a forest, or H must contain a component with negative (or zero) Euler characteristic, which means g is reducible. Both of these are contradictions. $\square$

4.5 Uniqueness of graph representatives

It is clear that given an efficient graph map compatible with a trellis map, the trellis mapping type can be reconstructed from the graph map. The converse is also true; up to isomorphism there is only one efficient controlled graph map compatible with a given trellis mapping class.

Theorem 4.30 (Uniqueness of graph representative) *Let $[f; T]$ be an irreducible trellis type. Then, up to isomorphism, there is a unique efficient graph map $(g; G)$ compatible with $[f; T]$.*

We first give a verifiable criterion for an exact homotopy equivalence between two graphs to be homotopic to a homeomorphism.

Lemma 4.31 *Suppose (G_1, W_1) and (G_2, W_2) are controlled graphs, and p is an exact homotopy equivalence between G_1 and G_2 . Suppose there are strictly positive real functions l_1 and l_2 on the edges of G_1 and G_2 such that whenever α_1 and α_2 are tight paths joining points of W with $p \circ \alpha_1 \sim \alpha_2$ we have $l_1(\alpha_1) = l_2(\alpha_2)$. (We take the distance between a point of W and a control vertex to be half the length of the control edge.) Then p is homotopic to a homeomorphism from (G_1, W_1) to (G_2, W_2) .*

PROOF: It is sufficient to show that the universal covers $(\tilde{G}_1, \tilde{W}_1)$ and $(\tilde{G}_2, \tilde{W}_2)$ are homeomorphic. Since $\tilde{W}_1$ and $\tilde{W}_2$ are homeomorphic, and G_1 and G_2 are homotopy-equivalent, the sets $\tilde{W}_1$ and $\tilde{W}_2$ are homeomorphic, and we call them both $\tilde{W}$. The functions l_1 and l_2 lift naturally to $\tilde{G}_1$ and $\tilde{G}_2$. For any four points $\tilde{w}_1, \tilde{w}_2, \tilde{w}_3$ and $\tilde{w}_4$ in $\tilde{W}$, there are only two possibilities for the span; either it contains a valence-4 vertex or two valence-3 vertices. Let α_{ij} be the curve from w_i to w_j .




Figure 29: The span of four points of W in a simply connected region can either have a valence-4 vertex (a) or two valence-3 vertices (b).

Suppose the span of $\{w_1, w_2, w_3, w_4\}$ contains a valence-4 vertex v as shown in Figure 29(a). Then for $i = 1, 2$,

$$l_i(\alpha_{12}) + l(\alpha_{34}) = l_i(\alpha_{13}) + l_i(\alpha_{24}) = l_i(\alpha_{14}) + l_i(\alpha_{23}) = l_i(\gamma_1) + l_i(\gamma_2) + l_i(\gamma_3) + l_i(\gamma_4)$$

However, if the span of $G\{w_1, w_2, w_3, w_4\}$ contains a valence-3 vertices v_1 and v_2 as shown in Figure 29(b), we have

$$l_i(\alpha_{12}) + l(\alpha_{34}) = l_i(\gamma_1) + l_i(\gamma_2) + l_i(\gamma_3) + l_i(\gamma_4)$$

but

$$l_i(\alpha_{13}) + l_i(\alpha_{24}) = l_i(\alpha_{14}) + l_i(\alpha_{23}) = l_i(\gamma_1) + l_i(\gamma_2) + l_i(\gamma_3) + l_i(\gamma_4) + 2l_i(\gamma_5),$$

so

$$l_i(\alpha_{12}) + l(\alpha_{34}) \leq l_i(\alpha_{13}) + l_i(\alpha_{24}) = l_i(\alpha_{14}) + l_i(\alpha_{23})$$

Hence the universal covers are homeomorphic by a homeomorphism $\tilde{h}$ homotopic to $\tilde{p}$. Further, since the computations given above are equivariant, the homeomorphism $\tilde{h}$ is equivariant, so projects to a homeomorphism $h : (G_1, W_1) \longrightarrow (G_2, W_2)$. $\square$

The proof of the main theorem is as follows:

PROOF: Let $p : (g_1; G_1) \longrightarrow (g_2; G_2)$ be an exact homotopy equivalence with homotopy inverse q . Thus $q \circ p \sim id$, $p \circ q \sim id$ and $g_1 \sim q \circ g_2 \circ p$, taking homotopies relative to W . It is clear that we can choose p and q to be mutual inverses on the control and peripheral edges, so merely need to consider the expanding edges. We therefore make no distinctions between control edges in G and G_2 .

First, consider a tight edge-path γ_1 in G_1 which has endpoints in W but no interior points in W . Let γ_2 be the tight edge-path homotopic to $p(\gamma_1)$. Consider γ_1 and γ_2 as exact curves $(I, J_0) \longrightarrow (G, W)$, and take minimal iterates under g_1 and g_2 . Since G_1 and G_2 are graphs, these minimal iterates can each be represented by a single tight edge-path. Then $\gamma_1 \sim q \circ p(\gamma_1) \sim q(\gamma_2)$, so $p((g_1)_{\min}(\gamma_1)) \sim (g_2)_{\min}(\gamma_2)$. Taking further iterates gives $p((g_1)_{\min}^n(\gamma_1)) \sim (g_2)_{\min}^n(\gamma_2)$ for all n . Thus $(g_1)_{\min}^n(\gamma_1)$ and $(g_2)_{\min}^n(\gamma_2)$ contain the same number of control and peripheral edges for all n . Since by Lemma 4.29, this number must be positive for all edges for some n , the graphs G_1 and G_2 are homeomorphic by a graph map h homotopic to p by Lemma 4.31. This homeomorphism gives a topological conjugacy between g_1 and g_2 . $\square$

5 Computing a Graph Representative

We now give an algorithm for obtaining an optimal controlled graph map compatible with a given trellis mapping class. If the trellis mapping class has is reducible, the algorithm may instead find a reduction, but can then be used to compute a graph representative for the irreducible components. The moves of the algorithm are based on the algorithm of Bestvina and Handel [BH95], but the method is quite different. Instead of reducing the topological entropy at each step, the algorithm reduces a zeta function, which gives a more precise measure of the growth based on the control and peripheral edges. This results in an algorithm where we need only consider the local action of the graph map.

In the case of a surface mapping class, there are no control edges, and no optimal graph map. We use a slightly different algorithm to find an efficient graph map, and consider a zeta function based on the peripheral edges to show that the algorithm terminates. If there are no peripheral edges, we create temporary peripheral edges by puncturing at a periodic orbit.

5.1 Graph moves

We now describe the moves we need for the algorithm. Each move is an exact homotopy equivalence; in particular, the Euler characteristic and number of control edges are unchanged. The property of being surface embeddable need not be preserved in general; however, performing any move followed by all possible vertex and edge tightenings does preserve this property.

The moves of edge tightening and vertex homotopy do not affect the graph (G, W) , but change the map g to a new map g' .

Edge tightening Suppose e_0 be an edge of G with $g(e_0) = e_1 \dots e_i e_{i+1} \dots e_n$ and $e_{i+1} = \bar{e}_i$. Homotope g to g' with $g'(e_0) = e_1 \dots e_{i-1} e_{i+2} e_n$.




Figure 30: Tightening the edge e .

Vertex homotopy Let v be a vertex of G with incident edges $e_1, \dots, e_k$, and let β be an edge path with $\iota(\beta) = g(v)$. Homotope g to a new map g' such that $g'(e_i) = \bar{\beta}g(e_i)$ for $i = 1, \dots, k$, unless $g(e_i)$ can be written $\beta\epsilon_i$ for some edge-path ϵ_i , in which case we take $g'(e_i) = \epsilon_i$. We say this move homotopes v *across* the edge-path β .

The moves of edge collapsing and vertex splitting change a graph G to a graph G' .

Edge collapsing Suppose e_0 is an edge of G with initial vertex v and final vertex w such that $g(e_0) = \cdot$. Suppose the cyclic ordering of edges at v is $\triangleleft e_0 \triangleleft a_1 \triangleleft \dots \triangleleft a_j \triangleleft e_0 \triangleleft$ and the cyclic ordering of edges at w is $\triangleleft \bar{e}_0 \triangleleft b_1 \triangleleft \dots \triangleleft b_k \triangleleft \bar{e}_0 \triangleleft$. Let G' be the graph formed by removing the vertex w and the edge e_0 from G such that the cyclic ordering of edges at v is $\triangleleft a_1 \triangleleft \dots \triangleleft a_j \triangleleft b_1 \triangleleft \dots \triangleleft b_k \triangleleft a_1 \triangleleft$. Let g' be the graph map obtained by removing all occurrences of e_0 from the image $g(e)$ of any edge e .

Vertex splitting Suppose v is a vertex of G , and the cyclic ordering of edges at v is $\triangleleft a_1 \triangleleft \dots \triangleleft a_j \triangleleft b_1 \triangleleft \dots \triangleleft b_k \triangleleft$. Let G' be a graph formed by introducing a new vertex w and new edge e_0 to G such that the cyclic ordering of edges at v is $\triangleleft e_0 \triangleleft a_1 \triangleleft \dots \triangleleft a_j \triangleleft e_0 \triangleleft$ and the cyclic ordering of edges at w is $\triangleleft \bar{e}_0 \triangleleft b_1 \triangleleft \dots \triangleleft b_k \triangleleft \bar{e}_0 \triangleleft$. Let g' be the graph map obtained by taking $g'(e_0) = \cdot$, and by replacing every occurrence of an edge b_i for $1 \leq i \leq k$ in the image $g(e)$ with the edge-path $e_0 b_i$ (and every occurrence of $\bar{b}_i$ with $\bar{b}_i \bar{e}_0$).

Edge collapsing and vertex splitting are inverse operations.

The following moves can all be expressed in terms of the basic moves described above.




Figure 31: (a) Before and (b) after a vertex splitting.

Vertex tightening If v is a vertex of G with incident edges $e_1, \dots, e_k$ and there is an edge-path β such that $g(e_i) = \beta\epsilon_i$ for all edges e_i with initial vertex v , homotope v across the edge-path β . The resulting graph map g' has $g'(e_i) = \epsilon_i$ for all edges e_i with initial vertex v .

Collapsing an invariant forest If $H \subset G$ is a set of expanding edges with $g(H) \subset H$ such that every component of H is simply connected (i.e. H is a forest), collapse every edge of T to a point. This move can be realised as a combination of homotopies and edge collapsing (though at intermediate stages, the set of control edges may not be invariant). The move is useful since the topology and dynamics of an invariant forest are trivial, and are removed to preserve transitivity of the graph map.

Tidying Perform edge tightenings, vertex tightenings, and collapsing invariant forests until no further tightening or collapsing is possible, with the exception that we never collapse peripheral edges at this stage (even if they have a trivial image).

Valence-3 homotopy Suppose v is a valence-3 vertex with incident edges e_1, e_2 and e_3 such that $g(e_i) = \beta\epsilon_i$ for $i = 1, 2$ and $g(e_3) = \epsilon_3$. Homotope v across β and to give $g'(e_i) = \epsilon_i$ for $i = 1, 2$ and $g'(e_3) = \bar{\beta}\epsilon_3$.

Folding Suppose there are edges $e_1 \triangleleft e_2$ at a vertex v with $g(e_1) = \epsilon_0\epsilon_1$ and $g(e_2) = \epsilon_0\epsilon_2$. First split off e_1 and e_2 to a new vertex w with new edge e_0 from v to w . We then perform any possible edge tightenings; note that since e_0 maps to $\cdot$, we cannot tighten at w . This gives a graph map g' with $g'(e_0) = \cdot$, $g'(e_1) = \epsilon'_0\epsilon'_1$ and $g'(e_2) = \epsilon'_0\epsilon'_2$. Finally, we homotope at w across the edge-path ϵ'_0 to obtain a graph map g'' with $g''(e_0) = \epsilon_0$, $g''(e_1) = \epsilon_1$ and $g''(e_2) = \epsilon_2$.

The analysis of tightening moves is simple, since they always decrease the complexity of the graph map. Similarly, collapsing invariant forests also decreases the complexity. Performing the tidying operation leave a nice graph map to use as a starting point for the more complicated moves of vertex homotopy and edge splitting.

Definition 5.1 (Tight and tidy graph maps) . We say a graph map is *tight* if no edge or vertex tightenings are possible. A tight graph map is *tidy* if it has no invariant forest of expanding edges, and every valence-2 vertex is an endpoint of a control edge.




Figure 32: Folding edges a and b .

5.2 Effect of graph moves

We measure the complexity of a graph map by its *zeta functions*.

Definition 5.2 (Zeta function) Let g be a graph map of a graph G and H an invariant subgraph. The H -length of an edge path ϵ , denoted $l[H]\epsilon$ is the number of occurrences of an edge of H in ϵ . The H -norm of a graph map g is given by

$$|g|_H = \sum_{e \in E(G)} l[H]g(e)$$

and the *control zeta function* of g is

$$\zeta_{g;H}(t) = \sum_{n=0}^{\infty} |g^n|_H t^n.$$

It is straightforward to show that if every invariant subgraph of G contains an edge in H , then the logarithm of the asymptotic growth rate of $|g^n|_H$ equals $h_{top}(g)$. If R is the radius of convergence of $\zeta_{g;H}$, the also $h_{top}(g) = \log(1/R)$, where R is

Zeta functions are ordered by the ordering $\sum_{n=0}^{\infty} a_n t^n < \sum_{n=0}^{\infty} b_n t^n$ if for some k we have $a_k < b_k$, but $a_i = b_i$ for $i < k$. If $k \leq N$, we say $\sum_{n=0}^{\infty} a_n t^n <_N \sum_{n=0}^{\infty} b_n t^n$. Equivalently, $\zeta_1 < \zeta_2$ if there exists $\epsilon > 0$ such that $\zeta_1(t) < \zeta_2(t)$ for all $t \in (0, \epsilon)$, and $\zeta_1 < N\zeta_2$ if $\zeta_1 + at^n < \zeta_2$ for all $a \geq 0$ and $n > N$.

The following lemma gives some simple results on H -norms.

Lemma 5.3 *Let g be a graph map and H an invariant subgraph of G . Then*

1. *If g' is obtained from g by edge tightening or vertex tightening, then $|(g')^n| \leq |g^n|_H$.*
2. *If g' is obtained from g by collapsing an edge e , then $|(g')^n|_H < |g^n|_H$ if $e \in H$ and $|(g')^n|_H = |g^n|_H$ otherwise.*
3. *If g' is obtained from g by collapsing an invariant forest F , then $|(g')^n|_H < |g^n|_H$ if $F \cap H \neq \emptyset$, and $|(g')^n|_H = |g^n|_H$ otherwise.*
4. *If g' is obtained from g by a vertex splitting introducing an edge e to give a new graph G' , then $|(g')^n|_H = |g^n|_H$.*

Valence-3 homotopies are more complicated, but the following result shows that H -norms are still reduced.

Lemma 5.4 *Let g be a graph map with a valence-3 vertex v which has incident edges e_1, e_2 and e_3 such that $g(e_i) = \beta\epsilon_i$ for $i = 1, 2$ and $g(e_3) = \epsilon_3$, and that g' is obtained from g by homotoping v across β . Suppose further that $g^{n-2}(\beta)$ does not contain any edges of H , but $g^{n-1}(\beta)$ does. Then*

1. *For any $k < n$ and any edge e of G , $l[H](g')^k(e) = l[H]g^k(e)$, and $|(g')^k|_H = |g^k|_H$.*
2. *$|(g')^n|_H < |g^n|_H$*

PROOF: Since $g^k(\beta)$ does not contain any control edges for $k < n - 1$, so $|g^k(\beta)|_H = 0$ for these values of k . For part 1, we use induction on k . The result is trivial for $k = 0$. Assume the result is true for some $k < n$. Then

$$\begin{aligned}
|(g')^{k+1}(e_i)|_H &= |(g')^k(\epsilon_i)|_H = |g^k(\epsilon_i)|_H \text{ (by the inductive hypothesis)} \\
&= |g^k(\beta)g^k(\epsilon_i)|_H - |g^k(\beta)|_H = |g^k(\beta\epsilon_i)|_H - |g^k(\beta)|_H \\
&= |g^{k+1}(e_i)|_H - |g^k(\beta)|_H \text{ for } i = 1, 2 \\
|(g')^{k+1}(e_3)|_H &= |(g')^k(\beta\epsilon_3)|_H = |g^k(\beta\epsilon_3)|_H = |g^k(\beta)|_H + |g^k(\epsilon_3)|_H \\
&= |g^k(e_3)|_H + |g^k(\beta)|_H \\
|(g')^{k+1}(e_i)|_H &= |(g')^k(\epsilon_i)|_H = |g^k(\epsilon_i)|_H = |g^{k+1}(e_i)|_H \text{ otherwise.}
\end{aligned}$$

Clearly then, $|(g')^k|_H = \sum_{e \in G} |(g')^k(e)|_H = \sum_{e \in G} |g^k(e)|_H = |g^k|_H$ for $k < n$. Part 2 follows from the above calculations since $|(g')^n|_H = |g^n|_H - |g^{n-1}(\beta)|_H < |g^n|_H$. $\square$

The following lemma shows that if we can find a bound on the number of edges of a graph, we can control which H -norm of g decreases in a valence-3 homotopy.

Lemma 5.5 *Suppose g is a map of a graph G with invariant subgraph H , and there are no invariant subgraphs of g which do not contain an edge in H . Suppose g' is obtained from g by a valence-3 homotopy. Then there exists $n \leq |G| - |H|$ such that $|(g')^k|_H = |g^k|_H$ for $k < n$ and $|(g')^n|_H = |g^n|_H$. In particular, $\zeta_{g';H} <_{|G|} \zeta_{g;H}$.*

PROOF: Let K_0 be the subgraph containing the edges of β , and define K_i by $K_i = K_{i-1} \cup g(K_{i-1})$. If K_n contains an edge of H for some least n , then $g^n(\beta)$ contains an edge of H and we are done. If not, then K_n has more edges than K_{n-1} , and hence has at least $n + 1$ edges. Therefore, K_n contains an edge of H for some $n \leq |G| - |H|$, the number of edges in $G \setminus H$. $\square$

The following lemma is used to show that our algorithm terminates.

Lemma 5.6 *Suppose an algorithm consists of graph moves such that there exists $N \in \mathbb{N}$ such that at each stage $\zeta_{g';H} <_N \zeta_{g;H}$. Then the algorithm terminates.*

PROOF: We use induction on N . Clearly, the result is true for $N = 0$, since $|g^0|_H = |H|$ is a non-negative integer.

Suppose the result holds for $N - 1$. Any move either decreases $|g^N|_H$ or $|g^k|_H$ for $k \leq N - 1$, but since $|g^N|_H$ is a non-negative integer, there is no infinite sequence of moves only decreasing $|g^N|_H$. By the inductive hypothesis, there can only be finitely many moves for which $|g^k|_H$ with $k < N$ is decreased, and the result follows. $\square$

5.3 Algorithm to compute the graph representative

We now give the algorithm to compute a graph representative, or find a reduction. An algorithm of Bestvina and Handel for free groups [BH92] considers the reducible case as a single graph.

Algorithm 5.7 Suppose g is a controlled graph map. Perform the following moves, followed by tidying (noting that tidying does not affect the peripheral subgraph), until the graph map is optimal or there is an invariant subgraph of free edges which is not a forest.

1. Fold a bad turn at a vertex of valence greater than 3, or at a control vertex.
2. Homotope at a bad turn at a free vertex of valence 3.
3. If no other moves are possible, collapse any peripheral edges which have trivial image.

The proof of termination considers zeta functions of the graph map. We need to consider both the control zeta function $\zeta_{g;C}$ and the peripheral zeta function $\zeta_{g;P}$, and also the number of peripheral edges $|P|$. The standard procedure for dealing with the peripheral subgraph P is *absorbing*, which yields a graph map for which there is no invariant set which deformation-retracts onto P other than P itself. However, absorbing into the peripheral subgraph can be accomplished by the moves of Algorithm 5.7. A fundamental observation is that G has at most $|G| = 2|C| - 3\chi$ edges, where χ is the Euler characteristic, and hence any decrease in a zeta function must occur in at most $|G|$ coefficients.

Theorem 5.8 (Computation of the graph representative) *Algorithm 5.7 terminates at a reduction or an optimal graph map.*

PROOF: If there is an invariant subgraph which does not contain any control or peripheral edges and is not a forest, g is irreducible and the algorithm has found a reduction. Further, at any stage there are at most $|E| = 2|C| - 3\chi$ edges of the graph, which means that any reduction of a zeta function must occur at most in the $|E|$ th coefficient. Therefore, every move apart from collapsing in the peripheral subgraph either reduces the control zeta function, increases the number of peripheral edges, or decreases the peripheral zeta function.

After collapsing peripheral edges in step 3, the link of the peripheral subgraph must be invariant under ∂g . If there is no reduction, every expanding edge in this subgraph must eventually map to a control edge, or else it would be contained in an invariant subgraph of free edges with negative Euler characteristic. We may perform homotopies across pre-peripheral edges disjoint from P which do not affect the control zeta function, but cannot introduce new peripheral edges without homotoping across an edge in $\text{Lk}(P)$. At this homotopy, the control zeta function decreases, so there is no infinite sequence of moves which does not decrease the control zeta function by Lemma 5.6. A further application of Lemma 5.6 shows that there are at most finitely many moves which decrease the control

zeta function. Hence the algorithm either finds a reduction, or terminates at an optimal graph map. $\square$

If there are no control edges, we are in the realm of Nielsen-Thurston theory. This case cannot be treated directly by our algorithm, since there is no optimal graph map in the homotopy class. The proof of Theorem 5.8 fails since after collapsing peripheral edges, all edges in $\text{Lk}(P)$ map to cover the entire graph, which is itself an invariant subgraph of negative Euler characteristic which does not include any control edges. Instead we consider the entropy in addition to the peripheral zeta function in order to show that the algorithm finds an efficient graph representative.

Algorithm 5.9 Suppose g is a graph map compatible with a surface mapping class, and g has nonempty peripheral subgraph P .

1. Perform Algorithm 5.7, stopping after collapsing peripheral edges in step 3.
2. If the graph map is efficient or has a reduction, the algorithm terminates. Otherwise, repeat step 1.

Perform Algorithm 5.7, checking after step 3 whether the graph map is efficient, at which point the algorithm terminates.

If there are no peripheral edges, we need to artificially introduce peripheral edges by puncturing at a periodic orbit.

Algorithm 5.10 Suppose g is a map of a graph G with no peripheral curves. Perform the following moves until an efficient graph map or a reduction is found.

1. Let P be a periodic orbit of g . Puncture at points of P by introducing artificial peripheral loops at P .
2. Perform the Algorithm 5.9 for graph maps with peripheral loops.
3. Collapse the artificial peripheral loops at P and tidy. If the resulting graph map is efficient or has a reduction, the algorithm terminates. If the resulting graph map has peripheral loops, perform Algorithm 5.9. Otherwise, return to step 1

Theorem 5.11 *Algorithms 5.9 and 5.10 terminate at a reduction or an efficient graph map.*

PROOF: We first consider Algorithm 5.9. If g is not efficient, there is a turn in the link of P which is inefficient. Folding this turn decreases entropy. Although other moves may increase entropy, the graph map obtained by folding and homotoping can also be obtained by the moves of the Bestvina-Handel algorithm, so the entropy must strictly decrease. Since the entropy is bounded above, and the number of edges of g is bounded by -3χ , there are only finitely many possible values for the entropy. Hence Algorithm 5.9 terminates.

Algorithm 5.10 terminates for similar reasons, since the entropy is bounded, and must strictly decrease. $\square$

We now give an example of computing an optimal graph map using Algorithm 5.7.

Example 5.12

Consider the trellis type $[f; T]$ shown in Figure 33.




Figure 33: A trellis with a period-2 orbit.

There are eleven stable segments, and the surface has two punctures, p and q , which are permuted by the diffeomorphism f .




Figure 34: An initial compatible graph

An initial compatible graph is shown in Figure 34. The edge-loop p_0p_1 surrounds the puncture p , and the edge-loop $q_0q_1q_2q_3$ surrounds the puncture q . The control edges map under g as follows:

$$z_0, z_1, z_2 \mapsto z_0; \quad z_3 \mapsto z_1; \quad z_4 \mapsto z_2; \quad z_5 \mapsto z_3; \quad z_6, z_7, z_8, z_9, z_{10} \mapsto z_4$$

We compute g on the expanding edges, which map:

$$\begin{aligned} p_0 &\mapsto p_0 \bar{z}_{10} c z_6 q_0; & p_1 &\mapsto q_1 q_2 q_3 \bar{z}_6 \bar{c} z_{10} \bar{p}_0; \\ q_0 &\mapsto q_1; & q_1 &\mapsto q_2 q_3 \bar{z}_6 \bar{c} z_{10} p_1; & q_2 &\mapsto p_0 \bar{z}_{10} c z_6 \bar{q}_3 \bar{q}_2 \bar{q}_1; & q_3 &\mapsto \cdot; \\ c &\mapsto \cdot; & d &\mapsto p_0 \bar{z}_{10} c z_6 \bar{q}_3 \bar{q}_2 \bar{q}_1; & e_0 &\mapsto e_0; & e_1 &\mapsto e_0; & e_2 &\mapsto e_1. \end{aligned}$$

Note that the edges p_i and q_i are not peripheral edges, since they do not form an invariant set. However the edge-loop p_0p_1 maps to the edge loop $p_0 \bar{z}_{10} c z_6 q_0 q_1 q_2 q_3 \bar{z}_6 \bar{c} z_{10} \bar{p}_0$ which is freely homotopic to $q_0 q_1 q_2 q_3$.

The edges q_3, c, e_1, e_2 and e_3 form an invariant forest which does not contain any control edges. Each of these edges can be collapsed to a point to give the graph in Figure 35.




Figure 35: Graph obtained by collapsing the invariant forest.

The expanding edges now map

$$\begin{aligned} p_0 &\mapsto p_0 \bar{z}_{10} z_6 q_0; & p_1 &\mapsto q_1 q_2 \bar{z}_6 z_{10} \bar{p}_0; \\ q_0 &\mapsto q_1; & q_1 &\mapsto q_2 \bar{z}_6 z_{10} p_1; & q_2, d &\mapsto p_0 \bar{z}_{10} z_6 \bar{q}_2 \bar{q}_1. \end{aligned}$$




Figure 36: Essential graph map.

The edge d has no preimage, so is not part of the essential graph representative. We can remove d , and also z_1 , z_3 , z_5 and z_8 to obtain the graph map shown in Figure 36. The control zeta function is $\zeta_{g;C}(t) = 5 + 13t + 25t^2 + \dots$.




Figure 37: Graph obtained after folding

Performing five folding operations introduced edges a_0 , a_1 , b_0 , b_1 and b_4 , as shown in Figure 37. Before tightening, the graph map is

$$\begin{aligned} p_0 &\mapsto a_0 p_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0 q_0 b_1; & p_1 &\mapsto \bar{b}_1 q_1 \bar{b}_4 b_4 q_2 b_0 \bar{z}_6 z_{10} a_1 \bar{p}_0 \bar{a}_0; \\ q_0 &\mapsto \bar{b}_1 q_1 \bar{b}_4; & q_1 &\mapsto b_4 q_2 b_0 \bar{z}_6 z_{10} a_1 p_1 \bar{a}_0; & q_2 &\mapsto a_0 p_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0 \bar{q}_2 \bar{b}_4 b_4 \bar{q}_1 b_1. \end{aligned}$$

On performing tightenings and valence-3 homotopies, we obtain the graph map

$$\begin{aligned} p_0 &\mapsto q_0; & p_1 &\mapsto q_1 q_2; & q_0 &\mapsto \cdot; & q_1 &\mapsto q_2 b_0 \bar{z}_6 z_{10} a_1 p_1; & q_2 &\mapsto p_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0 \bar{q}_2; \\ a_0 &\mapsto a_0 p_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0; & a_1 &\mapsto \bar{b}_1; & b_0 &\mapsto \bar{q}_1 b_1; & b_1 &\mapsto \bar{b}_4; & b_4 &\mapsto a_0. \end{aligned}$$

Collapsing q_0 and then p_0 gives the graph map shown in Figure 38, with

$$\begin{aligned} p_1 &\mapsto q_1 q_2; & q_1 &\mapsto q_2 b_0 \bar{z}_6 z_{10} a_1 p_1; & q_2 &\mapsto \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0 \bar{q}_2; \\ a_0 &\mapsto a_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0; & a_1 &\mapsto \bar{b}_1; & b_0 &\mapsto \bar{q}_1 b_1; & b_1 &\mapsto \bar{b}_4; & b_4 &\mapsto a_0. \end{aligned}$$




Figure 38: Graph obtained after collapsing q_0 and p_0 .




Figure 39: Graph obtained after folding q_1 and q_2 .

and control zeta function $\zeta_C(t) = 5 + 11t + 19t^2 + \dots$.

Folding q_1 and $\bar{q}_2$ to form a new edge b_2 gives the graph map shown in Figure 39, with

$$\begin{aligned} p_1 &\mapsto b_2 q_1 q_2 \bar{b}_2; & q_1 &\mapsto p_1; & q_2 &\mapsto \cdot; \\ a_0 &\mapsto a_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0; & a_1 &\mapsto \bar{b}_1; & b_0 &\mapsto \bar{q}_1 \bar{b}_2 b_1; & b_1 &\mapsto \bar{b}_4; & b_2 &\mapsto q_2 \bar{b}_2 b_0 \bar{z}_6 z_{10} a_1; & b_4 &\mapsto a_0. \end{aligned}$$




Figure 40: Graph obtained after collapsing q_2 .

Since now q_2 has trivial image, we can collapse q_2 to obtain the graph map shown in Figure 40, with

$$\begin{aligned} p_1 &\mapsto b_2 q_1 \bar{b}_2; & q_1 &\mapsto p_1; \\ a_0 &\mapsto a_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0; & a_1 &\mapsto \bar{b}_1; & b_0 &\mapsto \bar{q}_1 \bar{b}_2 b_1; & b_1 &\mapsto \bar{b}_4; & b_2 &\mapsto \bar{b}_2 b_0 \bar{z}_6 z_{10} a_1; & b_4 &\mapsto a_0. \end{aligned}$$

and control zeta function $\zeta_C(t) = 5 + 9t + 17t^2 + \dots$.

Finally, we fold up p_1 and q_1 to give new edges a_2 and b_3 as shown in Figure 41. The graph map is now

$$\begin{aligned} p_1 &\mapsto q_1; & q_1 &\mapsto p_1; \\ a_0 &\mapsto a_0 \bar{a}_1 \bar{z}_{10} z_6 \bar{b}_0; & a_1 &\mapsto \bar{b}_1; & a_2 &\mapsto b_3 \bar{b}_2 \\ b_0 &\mapsto \bar{b}_3 \bar{q}_1 b_3 \bar{b}_2 b_1; & b_1 &\mapsto \bar{b}_4; & b_2 &\mapsto \bar{b}_2 b_0 \bar{z}_6 z_{10} a_1; & b_3 &\mapsto a_2; & b_4 &\mapsto a_0. \end{aligned}$$

which is optimal. The control zeta function is $\zeta_{g;C}(t) = 5 + 9t + 15t^2 + \dots$. The edges p_1 and q_1 are peripheral edges for the graph map.




Figure 41: The optimal graph representative.

5.4 Algorithm for reduction

We finally give algorithms for performing reductions. We give two algorithms, one for separating reductions, and one for non-separating reductions. These algorithms are modelled on those of Keil [Kei97], but are simpler to apply in practice, since they do not require 2-simplices. Since reduction is not the main focus of this paper, and the algorithms do not introduce any significant new ideas, we provide only a brief analysis.

Algorithm 5.13 (Separating reduction) Let g be a controlled graph map with an invariant subgraph H consisting of n components, all of which have negative Euler characteristic and contain no control edges. Without loss of generality, we can assume the components of H are cyclically permuted by g .

1. Let p_1 be an edge in H such that the preceding edge in the cyclic order of vertices at $\iota(p_1)$ does not lie in H . Let π be the edge path p_1 .
2. Let p_k be the last edge of π , with $v_k = \iota(\bar{p}_k)$, and let p_{k+1} be next edge in the cyclic order of edges at v which lies in H .
 - If v_k is not the initial vertex of an edge in π , adjoin p_{k+1} to the end of π and return to step 2.
 - If p_{k+1} lies in π , this step terminates.

Otherwise, split off all edges from $\bar{p}_k$ to p_{k+1} inclusive.

- If $\bar{p}_{k+1}$ is not a directed edge in π , adjoin p_{k+1} to the end of π and return to step 2.
 - Otherwise, $\bar{p}_{k+1} = p_i$ in π . Homotope so that all edges at the new vertex v'_k so that $g'(p_{k+1}) = \cdot$, collapse p_{k+1} and return to step 2.
3. Let P be the frontier loop of H .
 4. Restrict g to a map g' of $G' = G \setminus H \cup P$. Tighten this restricted map.
 5. The output of the algorithm is a the pair of controlled graph maps $g^n|_K$, where K is a component of H , and g' on G' .

This algorithm works since once we have a situation where the frontier of H consists of simple closed curves, tightening g restricted to $G' = G \setminus H \cup P$ gives a graph map g' of G' by virtue of the fact that g is embeddable and acts as an automorphism of the fundamental group $\pi_1(G)$.

Algorithm 5.14 (Non-separating reduction)

1. Homotope to collapse all edges which eventually map into H .
2. Tighten g restricted to H , homotoping vertices if necessary.
3. Consider a component of H with vertices $v_0, v_1 \dots v_{k-1}, v_k = v_0$ and edges $p_1, \dots p_k$ with $\iota(p_i) = v_{i-1}$ and $\iota(\bar{p}_i) = v_i$. Create two new components of G , one with vertices v_i^+ and edges p_i^+ from v_{i-1}^+ to v_i^+ and one with vertices v_i^- and edges p_i^- from v_i^- to v_{i-1}^- . Move all the edges at vertex v_i in the interval $(\bar{p}_{i-1}, p_i)$ to the interval $(\bar{p}_{i-1}^+, p_i^+)$ at v_i^+ , and the edges in the interval $(p_i, \bar{p}_{i-1})$ to the interval $(\bar{p}_i^-, p_{i-1}^-)$ at v_i^- .
4. Let p be an edge in H , and suppose g is orientation-preserving. If $g(p) = q$ is a positively oriented edge, take $g'(p^+) = q^+$ and $g'(p^-) = q^-$, and if $g(p) = \bar{q}$, take $g'(p^+) = q^-$ and $g'(p^-) = q^+$. If instead g is orientation-reversing, take $g'(p^+) = \bar{q}^-$ and $g'(p^-) = \bar{q}^+$ if $g(p) = q$, and $g'(p^+) = \bar{q}^+$ and $g'(p^-) = \bar{q}^-$ if $g(p) = \bar{q}$.
5. Let e be an edge not in H with $g(e) = \epsilon\pi \dots$ where ϵ is an edge path with final edge e_k not in H , and $\pi = p_1 \dots p_j$ is an edge path in H . If $\iota(\bar{e}_k) = v^+$ for some vertex, take $g'(e) = \epsilon\pi^+ \dots = \epsilon p_1^+ p_2^+ \dots p_j^+ \dots$, and if $\iota(\bar{e}_k) = v^-$, take $g'(e) = \epsilon \bar{p}_1^- \bar{p}_2^- \dots \bar{p}_j^- \dots$.
6. H is not connected to the new graph G' and can be removed.
7. The output of the algorithm is a the controlled graph map g' of G' , together with an identification of pairs of peripheral loops $P^\pm$.

This algorithm works since the construction of the splitting of H into two copies of itself keeps track of the connections to the rest of the graph, and this allows us to easily construct a continuous graph map g' .

After step 1, all edges not in H map to at least one edge in H . After step 2, the edges in H are permuted; in particular, g is a homeomorphism on H . The final graph map is continuous since the separate definitions of g' on H' and $G' \setminus H'$ match up by considering orientations. In particular, note that since g is a homeomorphism on H and is embeddable, we cannot have an edge path ϵ in $G \setminus H$ which enters a component of H on one side and leave on the other.

6 Symbolic Dynamics and Shadowing

In this section, we discuss ways of representing the dynamics of a chaotic map. There are two main methods, the standard technique of symbolic dynamics, and the more powerful notion of *global shadowing*, which allows us to relate one map to another which is well understood.

6.1 Symbolic dynamics

Definition 6.1 (Itinerary and code) Let f be a map of a space X , and $\{R_\lambda\}_{\lambda \in \Lambda}$ be a finite set of compact subsets of X . Let (x_i) be an orbit of f , so $f(x_{i-1}) = x_i$ for all i .

Then a bi-infinite sequence $\dots R_{\lambda_{-2}} R_{\lambda_{-1}} R_{\lambda_0} R_{\lambda_1} R_{\lambda_2} \dots$ is an *itinerary* for (x_i) if $x_i \in R_{\lambda_i}$ for all i . If x_0 is a point of X , an $x_i = f^i(x_0)$ for all $i \geq 0$, then the infinite sequence $R_{n_0} R_{n_1} R_{n_2} \dots$ is an itinerary for x_0 if $x_i \in R_{n_i}$ for all $i \geq 0$. If x_0 is a period- n point of f , then the word $R_{\lambda_0} R_{\lambda_1} \dots R_{\lambda_{n-1}}$ is a *code* for x_0 if $x_i \in R_{\lambda_i}$ for $0 \leq i < n$.

Note that if the sets R_λ do not cover X , an orbit may not have any itinerary (if for some i , x_i is not in R_λ for any λ), and may have more than one itinerary if the sets R_λ are not disjoint. Typically, the sets R_λ are chosen to have disjoint interiors and boundaries with measure zero, so that the set of points with more than one itinerary has measure zero.

If $f : (X, Y) \longrightarrow (X, Y)$ is a map of pairs, we typically take the sets R_λ to have boundary in Y , and call such sets *regions*. If $(f; T)$ is a trellis map, we take itineraries of $\mathcal{C}f$ using sets R_λ which are regions of $(\mathcal{C}^U T, T^S)$. The itineraries of orbits of $\mathcal{C}f$ then give itineraries of orbits of f . One major advantage of trellis maps over other maps of surfaces is that the regions are specified in a natural way.

Definition 6.2 (Shift space) The *shift space* $\Sigma_\Lambda(f)$ of a map f on the regions $\{R_\lambda : \lambda \in \Lambda\}$ is the subset of $\{R_\lambda : \lambda \in \Lambda\}^\mathbb{N}$ consisting of itineraries of points under f .

The following result shows how the symbolic dynamics of the graph representative of a trellis mapping class relates to the dynamics of a trellis map. The proof is given in [Col99].

Theorem 6.3 *Let $(g; G, W)$ be the graph representative of an irreducible trellis mapping class $([f]; T)$, with G embedded as a subset of $\mathcal{C}^U T$. Then for every orbit (y_i) of g , there is an orbit (x_i) of f with the same itinerary as (y_i) . Further, if every region of T is simply-connected, then there is at most one orbit of g with a given itinerary (apart from orbits which enter the control set C).*

6.2 Global shadowing

Global shadowing is an equivalence relation on orbits of a system which gives a condition under which two orbits can be considered to be close to each other over infinite time intervals. The classical definition was introduced by Katok, and is given in terms of lifts of the map to the universal cover.

Definition 6.4 Let X be a Riemannian manifold with metric g and distance function d , and $f : X \longrightarrow X$ a continuous map. Let $\tilde{X}$ be the universal cover of X , let $\tilde{d}$ be the equivariant distance function obtained from the equivariant metric $\tilde{g}$ covering g , and let $\tilde{f} : \tilde{X} \longrightarrow \tilde{X}$ be a lift of f . Then orbits (x_i) and (y_i) *globally shadow* each other if they lift to orbits $(\tilde{x}_i)$ and $(\tilde{y}_i)$ of $\tilde{f}$ such that $\{\tilde{d}(\tilde{x}_i, \tilde{y}_i)\}$ is bounded.

This definition is independent of the metric used, and can be extended to certain equivariant metrics which are non-Riemannian.

As the concept is topological in spirit, it would be useful to have a purely topological classification. We will also be concerned with spaces which are topological pairs, and seek

a definition which extends naturally to this setting. We would also like to talk about arbitrary sets of orbits as well as just pairs of orbits. As a frequently-recurring theme in this work is that of homotopy classes of curves, a definition in terms of this formalism, rather than that of universal covers will also be useful. The basic idea of our global shadowing relation is given below. It relies on the definition of sets of curves of *bounded lengths*.

Definition 6.5 (Global shadowing) Orbits (x_i) and (y_i) of a map $f : (X, Y) \longrightarrow (X, Y)$ are said to *globally shadow* each other if there are sets $J_i \subset I$ such that $J_i \subset J_{i+1}$ and exact curves $\alpha_i : (I, J_i) \longrightarrow (X, Y)$ from x_i to y_i such that $f \circ \alpha_i$ is homotopic relative to endpoints to α_{i+1} via curves $(I, J_{i+1}) \longrightarrow (X, Y)$, and the curves α_i have *bounded lengths*. The curves α_i are called *relating curves*.

Similarly, (x_i) and (y_i) *forwards shadow* each other if there are shadowing curves for $i \geq 0$, and *backwards shadow* each other if there are shadowing curves for $i \leq 0$.

Note that we really need to allow the J_i to be an increasing sequence of sets; for example, we could take (x_i) and (y_i) to be orbits which start in X and map into Y .

There are many possible definitions of the term “bounded length,” and these mostly give identical definitions of global shadowing. In the setting of smooth curves on Riemannian manifolds, we can use the natural definition of the length of a smooth curve. In the setting of absolute neighbourhood retracts, we can use the following definition

Definition 6.6 (Bounded lengths) Let X be a compact ANR. A set of curves Γ has *bounded lengths* if for any closed ANR subset Z of X , there is a number $N(Z)$ such that each curve $\gamma \in \Gamma$ is homotopic to a curve α such that $\alpha(I) \cap Z$ has at most N components.

The following lemmas give some technical results which simplify the application of the global shadowing definition. The proofs are straightforward, so are omitted.

Lemma 6.7 *Let X be a manifold, and let Z be a set of disjoint hypersurfaces in X such that $X \setminus Z$ simply-connected. Suppose there is a number N such that for any $\gamma \in \Gamma$, γ has at most n intersections with Z . Then the curves in Γ have bounded lengths.*

Lemma 6.8 *Let Γ be a set of curves with the same endpoints which have bounded lengths. Then the set of isotopy classes $\{[\gamma] : \gamma \in \Gamma\}$ is finite.*

Lemma 6.9 *Let U be a contractible open subset of a space X . Then there is a set of curves Γ of bounded lengths contained in U such that for any two points x_0, x_1 in U , there is a curve in Γ from x_0 to x_1 .*

Given a homotopy between two different maps, we can also define what it means for an orbit of one map to globally shadow an orbit of the other map.

Definition 6.10 (Global shadowing for homotopic maps) Let $f_t : (X, Y) \longrightarrow (X, Y)$ be a homotopy from f_0 to f_1 , and $F : (X \times I, Y \times I) \longrightarrow (X \times I, Y \times I)$ be the fat homotopy of f_t defined by $F(x, t) = (f_t(x), t)$. Then an orbit (x_i) of f_0 and an

orbit (y_i) of f_1 are said to globally shadow each other under the homotopy f_t if the orbits $(x_i, 0)$ and $(y_i, 1)$ of F globally shadow each other under Definition 6.5.

6.3 Shadowing and Nielsen numbers

One of the main tools we have to relate the dynamics of maps with the same homotopy or isotopy type is Nielsen periodic point theory.

Definition 6.11 (Nielsen equivalence) Let f be a self-map of a topological pair (X, Y) . Period- n orbits (x_i) and (y_i) of f are *n-Nielsen equivalent* if there is a subset J of I and exact curves $\alpha_0, \dots, \alpha_{n-1} : (I, J) \longrightarrow (X, Y)$ from x_i to y_i such that $f \circ \alpha_i \sim \alpha_{i+1 \bmod n}$ (taking homotopies relative to endpoints).

Nielsen equivalence is the same as global shadowing for periodic orbits.

Lemma 6.12 *Periodic orbits (x_i) and (y_i) globally shadow each other if and only if they are n-Nielsen equivalent for some n .*

PROOF: Clearly, if (x_i) and (y_i) are Nielsen equivalent, they globally shadow each other. Conversely, if (x_i) and (y_i) globally shadow each other with shadowing curves (α_i) , then the possible homotopy classes for the α_i are bounded and the number of components of the J_i can be made to be bounded by suitable choices of the α_i . Hence, the J_i can be taken to be fixed, say $J_i = J$ for sufficiently large i . Then the homotopy classes of the α_i must eventually repeat, say $\alpha_{k+n} \sim \alpha_k$ for some $k \in \mathbb{Z}$ and $n > 0$. We can then take the curves to be $\beta_i = \alpha_{k+i}$ for $0 \leq i < n$. $\square$

It is easy to show [Col01] that Nielsen equivalence classes are closed in X , and hence have a well-defined fixed point index.

Definition 6.13 (Nielsen number) An n -Nielsen class is *essential* if its fixed-point index under f^n is nonzero. The n th Nielsen number of f , denoted $N_n(f)$ is the number of essential Nielsen classes of period- n orbits of f . If $R_{\lambda_0}, R_{\lambda_1}, \dots, R_{\lambda_{n-1}}$ is a code, then the its Nielsen number is the number of essential Nielsen classes with orbits of the code.

The most important property of exact homotopy equivalence is that it preserves the Nielsen numbers. Relative Nielsen theory can therefore be used to find periodic orbits and an entropy bound for a map of pairs. A complete exposition is given in [Col01].

The logarithm growth rate of the Nielsen numbers gives the *Nielsen entropy*.

Definition 6.14 (Nielsen entropy) The *Nielsen entropy* of a map f is the equal to the logarithm of the growth rate of the n th Nielsen numbers,

$$h_{niel}(f) = \limsup_{n \rightarrow \infty} \frac{\log N_n(f)}{n}.$$

6.4 Shadowing by trellis orbits

It is important to know how the dynamics of the graph representative models that of the trellis map. In Nielsen-Thurston theory, the orbits of the train-track map globally shadow those of the original surface homeomorphism. The results stated in Theorem 6.16 are analogous to the shadowing results of Nielsen-Thurston theory. We show that orbits of g are shadowed by orbits of f , and further, that periodic, asymptotic and biasymptotic orbits of g are shadowed by periodic, asymptotic and biasymptotic orbits of f .

We first need to clarify what it means for an orbit of g to be backward asymptotic to the control set C .

Definition 6.15 (Asymptotic graph orbits) An orbit (y_i) of g is *forward asymptotic* to C if $y_j \in C$ for some j . An orbit (y_i) of g is *backward asymptotic* to C if there exists $k \in \mathbb{Z}$ and exact curves $\mu_i : (I, \{0, 1\}) \rightarrow (G, W)$ such that μ_i begins in a periodic point of W , μ_i is homotopic in $\mathcal{C}T$ to a curve in T^U , $g \circ \mu_i \supset \mu_{i+1}$ for $i < k$ and $y_i \in \mu_i$.

Notice that if (y_i) is backward asymptotic to T^P , then the α -limit set of (y_i) lies in the control set C . The condition that the curves are homotopic to T^U is crucial; otherwise, we may have a periodic orbit of g which has α -limit set in C , but is not shadowed by an orbit of $\mathcal{C}f$ in T^U .

Theorem 6.16 (Shadowing by trellis orbits) Let $(f; T)$ be a trellis map and $(g; G, W)$ an efficient controlled graph map compatible with $(f; T)$. Then for any orbit (y_i) of g there is an orbit (x_i) of f which shadows (y_i) . Further,

1. If (y_i) is periodic, then (x_i) can be chosen to be periodic.
2. If (y_i) is forward asymptotic to C for some j , then (x_i) can be chosen in the stable set of T^P (so $x_j \in T^S$ for some j).
3. If (y_i) backward asymptotic to C , then (x_i) can be chosen in the unstable set of T^P .
4. If (y_i) is both forwards and backwards asymptotic to C , then (x_i) can be chosen to be biasymptotic to T^P .

Note that due to the asymmetry in our treatment of stable and unstable curves, part (2) and part (3) are genuinely different statements and require different proofs. As an immediate corollary, we have the following result.

Theorem 6.17 (Entropy of efficient graph maps) If g is an efficient controlled graph map compatible with $(f; T)$, then $h_{\text{top}}(g) \leq h_{\text{top}}(f)$.

We shall see in Section 7 that this entropy bound is sharp if T is well-formed, but otherwise need not be.

The remainder of this section is devoted to proving Theorem 6.16. We assume throughout that $(f; T)$ is a trellis map with graph representative $(g; G, W)$, and that (G, W) is embedded as a subset of $(\mathcal{C}^U T, T^S)$ by an embedding i . We let π be a deformation retract

$(\mathcal{C}^U T, T^S) \longrightarrow (G, W)$, and f_t be a homotopy from $\mathcal{C}f = f_0$ to $f_1 = i \circ g \circ \pi$ with fat homotopy F .

The following theorem is contained in [Col99]. It shows that periodic orbits of g are globally shadowed by an orbit of $\mathcal{C}f$, and forms the basis for all other shadowing results. The result is proved by showing that all Nielsen classes of g are essential.

Theorem 6.18 (Shadowing by periodic orbits) *Let (y_i) be a period- n orbits of g . Then there is a period- n orbit (x_i) of $\mathcal{C}f$ which is Nielsen equivalent to (y_i) .*

To show that all orbits of g are shadowed by orbits of f , we need to find a uniform bound for the lengths of shadowing curves in g .

Lemma 6.19 *If (x_i) and (y_i) are orbits of g which globally shadow each other, there are relating curves α_i from x_i to y_i such that the α_i lie in C , P or $\text{Pre-}P$.*

PROOF: Suppose the curves α_i are tight and do not cross any control edges. Then $\alpha_{i+1} = g \circ \alpha_i$ for all i , so the α_i are eventually periodic or disjoint. Since g has no invariant subgraphs apart from P , and g is expanding outside $\text{Pre-}P$, the α_i can only be eventually periodic if they lie in $\text{Pre-}P$.

If the α_i do cross C , they must eventually do so in periodic control edges. We can split the curves into pieces that cross C and pieces that do not. The pieces that do not must then lie in $\text{Pre-}P$. $\square$

Thus there is a uniform bound on the length of any relating curve in G . The following result shows that all orbits of g are shadowed by those of f .

Lemma 6.20 *Let (y_i) be an orbit of $(g; G, W)$. Then there is an orbit (x_i) of $(f; T)$ which shadows (y_i) .*

PROOF: If (y_i) is periodic, the result follows from the relative Nielsen theory. If not, suppose there are periodic orbits (y_i^j) such that for every i , $\lim_{j \rightarrow \infty} y_i^j = y_i$. We choose orbits (x_i^j) of f such that $(x_i^j, 0)$ shadows $(y_i^j, 1)$ under F . Since M is compact, by choosing a subsequence if necessary, we can assume the sequence x_0^j converges to a point x_0 as $j \rightarrow \infty$, and this means that x_i^j converges to $x_i = f_0^i(x_0)$ for all $i > 0$. Choosing further subsequences, we can assume further that x_i^j converges to a point x_i as $j \rightarrow \infty$ for any $i < 0$.

We therefore have an orbit (x_i) such that $x_i = \lim_{j \rightarrow \infty} x_i^j$ for all i . We now let π_i^j be the relating curves for $(x_i^j, 0) \sim (y_i^j, 1)$. Take an open cover $\mathcal{U}$ of $(\mathcal{C}^U T \times I, T^S \times I)$ by sets U such that the components of $U \setminus T^S$ are simply-connected. Let β_i^j and γ_i^j be curves from $(x_i, 0)$ to $(x_i^j, 0)$ and from $(y_i, 1)$ to $(y_i^j, 1)$ in a component of U and V respectively, which must exist for large enough j . Then $f_0 \circ \beta_i^j \sim \beta_{i+1}^j$ and $f_1 \circ \gamma_i^j \sim \gamma_{i+1}^j$ whenever j is sufficiently large.

Now let $\alpha_i^j = \beta_i^j \cdot \pi_i^j \cdot \gamma_i^j$. The curves α_0^j have bounded lengths, since they are joins from three sets of curves which each have bounded lengths. Therefore, there must be finitely

many homotopy classes $[\alpha_0^j]$, at least one of which must occur infinitely often. By taking an subsequence if necessary, we can assume $[\alpha_0^j]$ has the same value for all j . Now,

$$F \circ \alpha_i^j = F \circ \beta_i^j \cdot F \circ \pi_i^j \cdot F \circ \gamma_i^j \sim \beta_{i+1}^j \cdot \pi_{i+1}^j \cdot \gamma_{i+1}^j = \alpha_{i+1}^j$$

and since this must be true whenever α_i^j is defined, we have $[\alpha_i^j]$ is independent of j for fixed i . By taking subsequences if necessary, we can also ensure this is the case for i negative (or this follows automatically if F has a homotopy inverse). Hence, we have curves α_i from x_i to y_i of bounded lengths such that $F \circ \alpha_i \sim \alpha_{i+1}$. $\square$

6.5 Shadowing by stable and biasymptotic orbits

We now show that orbits of g which enter C are shadowed by orbits of f in $W^S(T^P)$, that orbits backward asymptotic to C are shadowed by orbits of f in $W^U(T^P)$, and that orbits of g which are biasymptotic to C are shadowed by orbits of f in $W^U(T^P) \cap W^S(T^P)$. These results have no corresponding statements in Nielsen-Thurston theory, since we cannot guarantee the existence of periodic saddle points in this case. We first prove an initial result which shows that orbits of g are contained in curves with endpoints in W which map over each other under g . If $\alpha_0 : (I, J) \longrightarrow (G, W)$ is a tight exact curve in G with endpoints in W , then we write $g(\alpha_0) \supset \alpha_1$ if $g(\alpha)$ contains α_1 as a sub-curve.

Lemma 6.21 *Let (y_i) be an orbit of g . Then there is a bi-infinite sequence of exact curves $\mu_i : (I, \{0, 1\}) \longrightarrow (G, W)$ such that $y_i \in \mu_i(I)$ and $g \circ \mu_i$ contains μ_{i+1} as a sub-curve.*

PROOF: First suppose that every component of $G \setminus W$ is homotopy-equivalent to a point or circle. Then every pair of points of W in the same region, there are two tight curves between them (for an homotopy circle) and one for a homotopy point. Let Γ be the set of all these curves. Let Σ_n , $n \geq 0$ be the set of bi-infinite sequences of elements of Γ such that $g(\gamma_i) \subset \gamma_{i+1}$ for $i \geq -n$. This is nonempty, since if $y_i \in \gamma_i$, then $y_{i+1} \in g(\gamma_i)$, and is compact since Γ is finite. Clearly $\Sigma_{n+1} \subset \Sigma_n$ so the Σ_n form a sequence of compact nested sets, hence $\Sigma = \bigcap_{n=0}^{\infty} \Sigma_n$ is nonempty. Then any sequence (μ_i) is a sequence in γ .

Now if g is irreducible optimal graph map, there exists N such that the complement of $g^{-N}(W)$ consists of homotopy points and circles. We consider g as a self-map of $(G, g^{-N}(W))$, and let (γ_n) be a sequence of curves found by the previous arguments. Then $g^N(\gamma_{i-N})$ is an exact curve $(I, \{0, 1\}) \longrightarrow (G, W)$ containing (y_i) , and maps across $g^N(\gamma_{i+1-N})$. Therefore taking $\mu_i = g^N(\gamma_{i-N})$ gives the required bi-infinite sequence of curves. $\square$

Lemma 6.22 *Let (y_i) be an orbit of g with $y_0 \in C$. Then there is an orbit (x_i) of f which globally shadows (y_i) with $x_0 \in T^S$.*

PROOF: Let (μ_i) be a sequence of exact curves $(I, \{0, 1\}) \longrightarrow (G, W)$ such that $g \circ \mu_i$ contains μ_{i+1} as a sub-curve. and $y_i \in \mu_i$. Let S_i be the stable segment containing y_i for $i \geq 0$; without loss of generality we assume $y_{-1} \notin T^S$.

62

and let μ_{i+1} be the tight curve in G which is exact homotopic to U_i , and so $y_{i+1} \in \mu_{i+1}$. Then the set $U = \bigcap_{n=0}^{\infty} f^{-n}(U_n)$ is a nonempty subset of T^U .




Figure 43: Construction of U_n .

We take $x_0 \in U$, and $(x_i) = f^i(x_0)$ for $i \neq 0$. A similar analysis to that of Lemma 6.22 shows that (x_i) globally shadows (y_i) . $\square$

Lemma 6.24 *Let (y_i) be an orbit of g such that for $y_k \in C$ and for $i \leq 0$ there are exact curves $\mu_i : (I, \{0, 1\}) \rightarrow (G, W)$ containing y_i such that $g(\mu_i) \supset \mu_{i+1}$, μ_i is unstable-parallel, has periodic initial vertex. Then there is an orbit (x_i) of f which shadows (y_i) such that $x_0 \in T^U$ and $x_k \in T^S$.*

PROOF: Without loss of generality we can assume that $y_k \in W$, but $y_{k-1} \notin C$. Take σ_i as in the proof of Lemma 6.23 for $i < k$, and let σ_k to be the stable segment S_k containing y_k . Construct U_i as in the proof of Lemma 6.23 for $i \leq k$. Then z_k is a point in $U_k \cap S_k$. We take $x_i = f^{i-k}(z_k)$. Then a similar analysis to that of Lemma 6.22 shows that (x_i) globally shadows (y_i) . Construct S_i for $i \geq n$ as in the proof of Lemma 6.22. $\square$

This completes the proof of Theorem 6.16

7 Entropy-Minimising Diffeomorphisms

In this section, we show that the entropy bounds obtained in Section 6 are sharp, at least for well-formed trellises. That is, the topological entropy of the graph representative g for a trellis type $[f; T]$ (which is the same as the Nielsen entropy $h_{niel}[f; T]$) is the infimum of the topological entropies of diffeomorphisms in the class. The main results of this section are summarised in the following theorem.

Theorem 7.1 (Existence of entropy minimising models) *Let $([f]; T)$ be a well-formed trellis mapping class. Then for any $\epsilon > 0$ there exists a diffeomorphism $\hat{f} \in ([f]; T)$ such that $h_{top}(\hat{f}) < h_{niel}[f; T] + \epsilon$. If there exists a diffeomorphism $\tilde{f} \in ([f]; T)$ such that any $\tilde{f}$ -extension of T is minimal, then there is a uniformly-hyperbolic diffeomorphism $\hat{f} \in ([f]; T)$ such that $h_{top}(\hat{f}) = h_{niel}[f; T]$. Further, if $([f]; T)$ is irreducible and $\tilde{f} \in ([f]; T)$ such that $h_{top}(\tilde{f}) = h_{niel}[f; T]$, then any $\tilde{f}$ -extension of T is minimal.*

We now give some examples which illustrate the hypotheses of the theorem. The following example shows that the hypothesis that the trellis be well-formed is necessary.

Example 7.2

The trellis mapping classes in Figure 44 are not well-formed.




Figure 44: Two ill-formed trellis mapping classes. The trellis in (a) has Nielsen entropy $\log 2$, whereas the trellis in (b), which is the time reversal of that in (a), has Nielsen entropy zero.

The Nielsen entropy of the trellis mapping class in (a) is equal to $\log 2$, so any diffeomorphism in the trellis mapping class must have topological entropy at least $\log 2$. Since the Smale horseshoe map has this trellis type, the topological entropy of the trellis type is exactly $\log 2$. The trellis mapping class of Figure 44(b) is conjugate to the time-reversal of the trellis mapping class in (a). Since the topological entropy of a diffeomorphism is the same as that of its inverse, any diffeomorphism in this trellis mapping class must have topological entropy at least $\log 2$. However, all the edges of the graph representative are control edges, so the Nielsen entropy is zero.

The above example illustrates that a trellis which is not well-formed may have Nielsen entropy strictly less than the topological entropy, and may even have different Nielsen entropy from its time-reversal. Even if a trellis mapping class is well-formed, it is not necessarily true that the Nielsen entropy is realised. A trivial example is of a well-formed trellis type $[f; T]$ for which $h_{niel}[f; T] = h_{top}[f; T]$ by for which every diffeomorphism in $[f; T]$ has topological entropy greater than $h_{niel}[f; T]$ is the planar trellis type with a single transverse homoclinic intersection. The Nielsen entropy of this trellis type is equal to zero, but every diffeomorphism with a transverse homoclinic point has strictly positive topological entropy. However, it is simple to construct trellis maps with topological entropy arbitrarily close to zero.

We now give a nontrivial example.

Example 7.3

A trellis type $[f; T]$ for which the Nielsen entropy is not realisable is shown in Figure 45. Consider the segment S . Taking backward minimal iterates of S eventually yields a segment lying in the region $R(Q)$ with endpoints in the segment $U_P(Q)$. Then, by the Lambda lemma, under any diffeomorphism f in the trellis mapping class, $f^{-n}(S)$ tends to the closed branch of $T^S(p)$ as $n \rightarrow -\infty$, so contains an intersection with T^U for some n , even though any *minimal* backward iterate of S has no intersections with T^U .




Figure 45: A trellis for which the Nielsen entropy is not realisable.

Similarly, $f^n(U)$ must intersect T^S for some n even though any minimal iterate does not. These observations can be used to show that the Nielsen entropy of the extension $[f; \widehat{T}]$ is greater than that of $[f; T]$, so $h_{top}(f) > h_{niel}[f; T]$. In general, the realisability of the entropy bound is closely related to the existence of a diffeomorphism for which every extension is a minimal extension.

In the case where this infimum is realised, we show how to construct a minimal-entropy uniformly-hyperbolic diffeomorphism in the trellis mapping class. Otherwise, we show, for any $\epsilon > 0$, how to construct a diffeomorphism whose entropy is within ϵ of the Nielsen entropy. The following example gives a trellis mapping class for which the Nielsen entropy cannot be realised.

7.1 Nielsen entropy of minimal trellises

Most of the procedures we use to construct diffeomorphisms in a given trellis mapping class rely on extending the original trellis and introducing new branches in a controlled way. The most important type of extension is a minimal extension, since, as noted in Section 3.4, we expect the Nielsen entropy of a minimal extension to be the same as that of the original trellis mapping class. We show this in Theorem 7.5, using results on the graph representative of a minimal stable supertrellis given in Lemma 7.4. We also consider methods for introducing new points of T^P without increasing the Nielsen entropy. The only difficulty here is on finding the correct initial segment of a branch; once this has been achieved, we can take minimal iterates in the usual way. Since the Nielsen entropy is computed via a graph representative, which induces an asymmetry between unstable and stable curves, we need to consider each separately. We consider initial unstable segments in Lemma 7.9, and initial unstable segments in Lemma 7.10. In Theorem 7.11 we apply these results to the cases we need to consider later.

Lemma 7.4 *Let $([f]; T)$ be an irreducible trellis mapping class with graph representative $(g; G, W)$, and let $([f]; \widehat{T})$ be a minimal stable supertrellis with graph representative $(\tilde{g}; \tilde{G}, \tilde{W})$. Then there is an embedding $i : (G, W) \hookrightarrow (\tilde{G}, \tilde{W})$ such that $i \circ g = \tilde{g} \circ i$, and i restricts to a homeomorphism between the invariant subgraphs $\bigcup_{i=0}^{\infty} g^i(G)$ and $\bigcup_{i=0}^{\infty} \tilde{g}^i(\tilde{G})$*

Further, if l is a length function on G such that $l(g(z)) = l(z)$ for any periodic control edge, $l(g(p)) = l(p)$ for any peripheral edge, and $\lambda l(e) \leq l(g(e)) < \lambda_\epsilon l(e)$ for any other edge, there is a length function $\tilde{l}$ on $\tilde{G}$ with the same properties such that $\tilde{l}(i(e)) = l(e)$ for any edge of G .

PROOF: First note that $(\tilde{G}; W)$ is exact homotopy equivalence to $(G; W)$, and this exact homotopy equivalence also gives an exact homotopy equivalence between $\tilde{g}$ considered as a map of $(\tilde{G}; W)$ and g . There is also a natural embedding $W \hookrightarrow \tilde{W}$ induced by the inclusion on the stable segments.

Let $\tilde{\alpha}$ be a path in $\tilde{G}$ with endpoints only in W and minimal intersections with $\tilde{W}$. Since $\tilde{f}$ is a minimal supertrellis, the path $\tilde{g}(\alpha)$ also does not back-track. Since any vertex of (G, W) is contained in the span of the control set C , this is enough to prove that i is homotopic to an embedding. Therefore all control edges of $\tilde{G}$ are either in the span of C , or in a forest which is attached to the span of C . Since all control edges of this forest map into C , and have no control edge preimages, all edges of $\tilde{G}$ map into the span of C under $\tilde{g}$.

Any edge of $\tilde{G}$ maps to an edge-path which is the embedding of an edge of G under $\tilde{g}$. We take the length function $\tilde{l}$ in the span of C to be given by $\tilde{l}(\tilde{e}) = l(g(\tilde{e})) * (l(e)/l(g(e)))$ where e is the edge of g containing $\tilde{e}$. We take the length of an edge $\tilde{e}$ not in the span of C to be $l(g(\tilde{e}))/\lambda_*$, where $\lambda < \lambda_* < \lambda_\epsilon$. This gives the required length function. $\square$

Note that a stable supertrellis which is not minimal may change the topology of the graph.

Theorem 7.5 (Nielsen entropy of minimal extensions) *Let $([f]; T)$ be a well-formed trellis mapping class. If $([\tilde{f}]; \tilde{T})$ is a minimal extension of $([f]; T)$, then $h_{niel}[\tilde{f}; \tilde{T}] = h_{niel}[f; T]$.*

PROOF: There exists n such that $\tilde{f}^{-n}(\tilde{T}^U \subset T^U$. Let $([\hat{f}]; \hat{T})$ be a minimal extension of $([\tilde{f}]; \tilde{T})$ such that $\hat{f}^{-n}(\hat{T}^U) \subset T^U$. Then $([\hat{f}]; \hat{T})$ is conjugate to the trellis mapping class $([\hat{f}]; (\hat{f}^{-n}(T^U); \hat{f}^{-n}(T^S)))$, which is a minimal stable extension of $([f]; T)$, so, by Lemma 7.4, $h_{niel}[f; T] = h_{niel}[\hat{f}; (\hat{f}^{-n}(T^U); \hat{f}^{-n}(T^S))]$. Hence

$$h_{niel}[f; T] \leq h_{niel}[\tilde{f}; \tilde{T}] \leq h_{niel}[\hat{f}; \hat{T}] = h_{niel}[\hat{f}; (\hat{f}^{-n}(\hat{T}^U); \hat{f}^{-n}(\hat{T}^S))] = h_{niel}[f; T],$$

yielding $h_{niel}[\tilde{f}; \tilde{T}] = h_{niel}[f; T]$ as required. $\square$

We now give an example of a trellis and a minimal stable iterate, together with the graph representatives.

Example 7.6

The minimal extension of the trellis map shown in Figure 46(a) is shown in Figure 46(b), together with the graph representatives.

The control edges for the original trellis are shown with shaded dots, and those of the extension with white dots. This clearly shows the embedding of the graph representative of the trellis in that of its minimal extension. In each case, the essential graph representative is the span of $\{z_0, z_2, z_4\}$.




Figure 46: (a) A trellis type with its graph representative, and (b) the minimal stable iterate.

We next state a simple result which shows that we can puncture at points of an essential periodic orbit without changing the Nielsen entropy.

Lemma 7.7 *Let $([f]; T)$ be a well-formed trellis mapping class with graph representative $(g; G, W)$. Suppose P is an essential periodic orbit of f which does not globally shadow T^S or ∂M . Let $([\tilde{f}]; \tilde{T})$ be the trellis mapping class obtained by puncturing M at the points of P . Then $h_{\text{niel}}[\tilde{f}; \tilde{T}] = h_{\text{niel}}[f; T]$.*

PROOF: Since P is an essential periodic orbit of f , it is shadowed by a periodic orbit of g . Replacing the periodic orbit of g shadowing P with a peripheral loop yields the graph representative $\tilde{g}$ of $([\tilde{f}]; \tilde{T})$, which clearly has the same topological entropy as that of g . $\square$

We now turn to supertrellises formed by introducing new branches. The next lemma applies to both stable and unstable supertrellises, since it deals only with minimal iterates of arcs and cross-cuts to T .

Lemma 7.8 *Let $([f]; T)$ be a trellis mapping class, and let A be a set of mutually disjoint embedded curves in $\mathcal{C}^U T$ with endpoints, but no interior points, in T^S . Additionally, if a point of A in an arc α lies in a periodic segment S of T^S , then α initially lies along an unstable segment of T^U . Suppose there is a minimal iterate B of A such that every arc of A is contained in an curve of B . Then there is a minimal unstable supertrellis $([\tilde{f}]; \tilde{T})$ of $([f]; T)$ such that $\tilde{T}^U = T^U \cup B$.*

Similarly, if A is a set of cross-cuts to T^U which are disjoint from T^S , and there is a minimal iterate B of A such that $A \subset B$, then there is a minimal unstable supertrellis $([\tilde{f}]; \tilde{T})$ of $([f]; T)$ such that $\tilde{T}^U = T^U \cup B$.

PROOF: Let $\beta_{i+1} \in f_{\min}[\alpha_i]$ and be disjoint from T^U . Without loss of generality, we can assume that $\beta_{i+1} \supset \alpha_{i+1}$ for all i . Then the curves $f \circ \alpha_i$ and β_{i+1} are isotopic by an

isotopy mapping the endpoints into $f(T^S)$. Further, since the endpoints of curves of α do not lie in segments of T^S containing points of T^P , or are points of T^P , we can take h_t to be fixed these segments. By the isotopy extension theorem A.2 this isotopy can be extended to an ambient isotopy h_t which is fixed on T^U and maps $f(T^S)$ into $f(T^S)$. Let $\tilde{f} = h_1 \circ f$, so T^U and T^S are still part of the unstable and stable manifolds of T^P for $\tilde{f}$. Then $\tilde{f}(T^U) = h_1(f(T^U)) \supset h_1(T^U) = T^U$, $\tilde{f}(T^S) = h_1(f(T^S)) = f(T^S) \subset T^S$ and $\tilde{f}(\alpha_i) = h_1(f(\alpha_i)) = \beta_{i+1} \supset \alpha_{i+1}$. Therefore $([\tilde{f}]; \tilde{T})$, where $\tilde{T} = (T^U \cup \bigcup \beta_i, T^S)$, is a well-formed trellis mapping class, and is a minimal supertrellis of $([f]; T)$ since the curves β are minimal iterates of the curves α . The statement for cross-cuts to T^U follows by reversing time. $\square$

We can use this lemma to show the existence of a minimal unstable supertrellis based on the graph map, and that the Nielsen entropy of this supertrellis is equal to the Nielsen entropy of the original trellis type.

Lemma 7.9 *Let $([f]; T)$ be a well-formed trellis mapping class with graph representative $(g; G, W)$. Let γ_i for $0 \leq i < n$ be a set of curves in G with endpoints only in such that $g \circ \gamma_i$ contains $\gamma_{i+1 \bmod n}$ as a sub-curve, and each γ_i lifts to an embedded curve α_i in $C^U T$ such that the α_i are disjoint. Suppose further that if an endpoint of some γ_i lies in a periodic point $w \in W$, then γ_i lies on the same side of W as a trivial branch of T . Then the collection $A = \{\alpha_i\}$ satisfies the conditions of Lemma 7.8, and the minimal supertrellis $([\tilde{f}]; \tilde{T})$ so obtained has $h_{\text{niel}}[\tilde{f}; \tilde{T}] = h_{\text{niel}}[f; T]$.*

PROOF: The curves $f \circ \alpha_i$ are homotopic to $\epsilon_{i+1} = g \circ \gamma_i$, so are homotopic to minimal iterates β_{i+1} of the α_i . By a further homotopy we can ensure α_i is contained in β_i , and so satisfies the hypotheses of Lemma 7.8. The graph representative $(\tilde{g}; \tilde{G}, \tilde{W})$ is given by cutting along the curves ϵ_i , so there is a projection $\pi : (\tilde{G}, \tilde{W}) \rightarrow (G, W)$ such that $\pi \circ \tilde{g} = g \circ \pi$. Since $\tilde{g}$ is optimal, this ensures $h_{\text{top}}(\tilde{g}) = h_{\text{top}}(g)$. $\square$

We have already shown that the Nielsen entropy of a minimal stable supertrellis is the same as that of the original trellis type. The following result shows how to construct minimal stable supertrellises.

Lemma 7.10 *Let $([f]; T)$ be a well-formed trellis mapping class with graph representative $(g; G, W)$. Suppose P is an essential periodic orbit of f which does not globally shadow T^S or ∂M . Then f is isotopic relative to P to a diffeomorphism $\tilde{f}$ with a trellis $\tilde{T}$ such that $([\tilde{f}]; \tilde{T})$ is a well-formed supertrellis of $([f]; T)$ such that P is contained in an attractor of $([\tilde{f}]; \tilde{T})$ and $h_{\text{niel}}[\tilde{f}; \tilde{T}] = h_{\text{niel}}[f; T]$.*

PROOF: Take cross-cuts A in the statement of Lemma 7.8 to have essential intersections with the edges of g containing P , giving trellises and controlled graphs shown in Figure 47. Then the supertrellis $([\tilde{f}]; \tilde{T})$ given by Lemma 7.8 is the required trellis mapping class. $\square$

We use these results to construct new branches at essential periodic orbits and trivial branches of a trellis. These cases are shown in Figure 48. These branches are may used to create new attractors and repellers, or subdivide a region into an attractor/repellor and




Figure 47: Trellises and graphs formed by introducing new stable curves edges. The in (a) formed by introducing a single stable curves does not have any attractors. The trellises in (b) and (c) formed by introducing one stable curve crossing each incident edge at a vertex do yield new attracting regions.




Figure 48: Stable supertrellises. In (a) we introduce an attractor, in (b) we bound an attractor, and in (c) we extend a trivial branch.

a collection of rectangles, as in Figure 48(a), to bound the end of a trivial branch in an attractor or repellor, as in (b), or to extend a trivial branch to a nontrivial branch, as in (c). These cases correspond to the three cases of Theorem 7.11

Theorem 7.11 *Let $([f]; T)$ be a well-formed trellis mapping class. Then:*

- (a). *If P is an essential periodic orbit of $([f]; T)$ which does not shadow T^P , there is are minimal supertrellises of $([f]; T)$ for which P is contained in a region which is attracting, repelling, or both.*
- (b). *If P is a periodic orbit at the ends of trivial branches of T^P , then there is a minimal supertrellis $([\tilde{f}]; \tilde{T})$ of $([f]; T)$ for which P is contained in a non-chaotic region of $([\tilde{f}]; \tilde{T})$.*
- (c). *If $T^{U/S}(p, b)$ is a trivial branch of T which lies in a chaotic region of T , then there is a minimal supertrellis $([\tilde{f}]; \tilde{T})$ of $([f]; T)$ for which $\tilde{T}^{U/S}(p, b)$ is contained in a nontrivial branch of T^P .*

All these statements follow from Lemma 7.9 and Lemma 7.10 by constructing the required curves or finding the right control edges.

7.2 Regular domains

Any diffeomorphism is conjugate to a linear hyperbolic diffeomorphism in a neighbourhood of a point of a periodic saddle point. We therefore expect a trellis map $([f]; T)$ to behave

in a fairly predictable way in a neighbourhood of a point of T^P . In particular, we know how a rectangle with sides parallel to the local stable and unstable foliations behaves. Unfortunately, for a given trellis mapping class, the neighbourhood of T^P on which we have hyperbolic behaviour may be arbitrarily small. To deal with this problem, we introduce the concept of a *regular domain*, which is a rectangular domain which behaves similarly to a sufficiently small rectangular neighbourhood of a periodic saddle point.

Definition 7.12 (Regular domain) Let $([f]; T)$ be a trellis mapping class. A rectangular domain D of T is a *regular domain* for a period- n quadrant Q of T if D has sides $T^U[p, q^u] \subset f^{-n}(T^U)$, $T^S[p, q^s] \subset f^n(T^S)$, $T^U[q^s, r]$ and $T^S[q^u, r]$, such that $f^n(T^U[p, q^u]) \cap D = T^U[p, q^u]$, $f^{-n}(T^S[p, q^s]) \cap D = T^S[p, q^s]$, and $f^{-n}(T^U[q^s, r]) \cap D = f^n(T^S[q^u, r]) \cap D = \emptyset$.

The sides $T^U[p, q^u]$ and $T^S[p, q^s]$ are called *adjacent sides* of D , and the sides $T^U[q^s, r]$ and $T^S[q^u, r]$ are *opposite sides*. Note that the definition is given purely in terms of the topology of the trellis and the mapping of its vertices; no extensions are needed.




Figure 49: The domain D is a regular domain for Q .

A regular domain is shown in Figure 49. We shall always denote the vertices of a regular domain by p , q^u , q^s and r as shown. We now give a number of elementary properties of a regular domain. The first two lemmas are trivial.

Lemma 7.13 Let $([\hat{f}]; \hat{T})$ be an extension of $([f]; T)$. Then if D is a regular domain for a quadrant Q in $([f]; T)$, then D is also a regular domain for Q in $([\hat{f}]; \hat{T})$.

Lemma 7.14 Let D be a regular domain of Q , and suppose that both $f^{-(n+1)}(T^S[p, q^s])$ and $f^{-1}(T^S[q^u, r])$ are subsets of T^S . Then $f^{-1}(D)$ is a regular domain for $f^{-1}(Q)$.

Lemma 7.15 Let D be a regular domain for Q , and $T^S[\tilde{q}^u, \tilde{r}] \subset D$ be a curve with endpoints $\tilde{q}^u \in T^U(p, q^u)$ and $\tilde{r} \in T^U(q^s, r)$. Then the rectangular domain $\tilde{D}$ with vertices at $\{p, \tilde{q}^u, q^s, \tilde{r}\}$ is a regular domain for Q . Similarly, if $T^U[\tilde{q}^s, \tilde{r}] \subset D$ is a curve with endpoints $\tilde{q}^s \in T^S(p, q^s)$ and $\tilde{r} \in T^S(q^u, r)$, then the rectangular domain vertices at $\{p, q^u, \tilde{q}^s, \tilde{r}\}$ is a regular domain for Q .

PROOF: The only nontrivial step is to show $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap \tilde{D} = \emptyset$. First, note that $f^n(\tilde{q}^u) \in T^U(\tilde{q}^u, f^n(q^u))$, so $f^n(\tilde{q}^u) \notin \tilde{D}$. Further, $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap T^U[p, \tilde{q}^u] = f^n(T^S[\tilde{q}^u, \tilde{r}] \cap f^{-n}(T^U[p, \tilde{q}^u])) = f^n(\emptyset) = \emptyset$ and $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap T^U[p, q^u] = f^n(T^S[\tilde{q}^u, \tilde{r}] \cap f^{-n}(T^U[p, q^u])) \subset f^n(D \cap f^{-n}(T^U[p, q^u])) = \emptyset$, so $f^n(T^S[\tilde{q}^u, \tilde{r}])$ does

not intersect the unstable boundary of $\tilde{D}$. Additionally, $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap T^S[p, q^s] = f^n(T^S[\tilde{q}^u, \tilde{r}] \cap f^{-n}(T^S[p, q^s])) \subset f^n(T^S[\tilde{q}^u, \tilde{r}] \cap T^S[p, q^s]) = \emptyset$. It remains to show that $f^n(T^S[\tilde{q}^u, \tilde{r}])$ does not intersect itself. We have already seen that $f^n(T^S[\tilde{q}^u, \tilde{r}])$ does not contain $\tilde{q}^u$ or $\tilde{r}$. Further, $f^n(\{\tilde{q}^u\}) \cap T^S[\tilde{q}^u, \tilde{r}] \subset f^n(T^U(p, q^u)) \cap T^S[\tilde{q}^u, \tilde{r}] = \{\tilde{q}\}$, but clearly $f^n(\tilde{q}^u) \neq \tilde{q}^u$, so $f^n(\tilde{q}^u) \notin T^S[\tilde{q}^u, \tilde{r}]$. Since $T^S[\tilde{q}^u, \tilde{r}]$ is an interval, this is enough to show that $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap T^S[\tilde{q}^u, \tilde{r}] = \emptyset$. Hence $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap \partial\tilde{D} = \emptyset$, so $f^n(T^S[\tilde{q}^u, \tilde{r}]) \cap \tilde{D} = \emptyset$. A similar analysis proves the statement for a curve $T^U[\tilde{q}^s, \tilde{r}]$. $\square$




Figure 50: $\tilde{D}$ is a regular domain if D is a regular domain.

Two applications of Lemma 7.15 show that the region $\tilde{D} \subset D$ shown in Figure 50 is a regular domain for Q if D is a regular domain.

One of the most important properties of a regular domain is that iterates of the sides opposite Q must cross each other.

Lemma 7.16 *Suppose D is a regular domain for a period- n quadrant Q , and suppose $f^{-n}(T^S[q^u, r]) \subset T^S$. Then $f^{-n}(T^S[q^u, r])$ contains a subinterval of the form $T^S[\tilde{q}^u, \tilde{r}] \subset D$ such that $\tilde{q}^u = f^{-n}(q^u) \in T^U(p, q^u)$ and $\tilde{r} \in T^S(q^s, r)$.*

PROOF: Since $f^n(Q) = Q$, f^n is orientation-preserving, so the orientation of the intersection at $f^{-n}(q^u)$ is the same as that at q^u . Further, $f^{-n}(T^S[q^u, r]) \cap T^U[p, q^u] = f^{-n}(T^S[q^u, r] \cap f^n(T^U[p, q^u])) = f^{-n}(T^S[q^u, r] \cap T^U[p, q^u]) = \{f^{-n}(q^u)\}$, so $f^{-n}(T^S[q^u, r])$ does not cross $T^U[p, q^u]$. Finally, $\{f^{-n}(r)\} \cap D \subset f^{-n}(T^U[q^s, r]) \cap D = \emptyset$, so $f^{-n}(r) \notin D$. Therefore, $f^{-n}(T^S[q^u, r])$ intersects $T^U[q^s, r]$, and we let $\tilde{r}$ be this first intersection. $\square$

Notice that Lemma 7.16 is enough to show that if D is a regular domain for a trellis mapping class $([f]; T)$, and $([f]; T)$ is an extension of $([f]; T)$ with $\hat{f}^{-n}(T^S[q^u, r]) \subset \hat{T}^S$, then the biasymptotic orbit through $\tilde{r}$ is forced by $([f]; T)$.

In Figure 50 we show a regular domain D together with $f^{-n}(T^S[q^u, r])$. Notice that $f^{-n}(T^S[q^u, r])$ may cross $T^U[q^s, r]$ several times, but must cross at least once since $f^{-n}(r) \notin D$.

The most important case of a regular domain for a quadrant Q is when D is a region, in which case we have a *regular region* for Q . It is also useful to consider the less restrictive case for which the interior of D does not intersect T^U , or in other words, if $D \cap T^U \subset \partial D$. In this case, there is a unique homotopy class of exact curves $\alpha_D : (I, J) \rightarrow (\mathcal{C}^U T, T^S)$ such that $\alpha_D(I) \subset D$, $\alpha_D(0) \in T^U[p, q^s]$, $\alpha_D(1) \in T^U[q^u, r]$, and α_D has minimal intersections




Figure 51: The backward iterate of $T^S[q^u, r]$ intersects $T^U[q^s, r]$ but does not cross $T^U[p, q^u]$.

with T^S . It is an important but trivial observation that if $D \cap T^U = \partial D^U$ and if $\hat{T}$ is an extension of T with $\hat{T}^U = T^U$, then $D \cap \hat{T}^U = \partial D^U$. The following lemma shows that any curve in such a regular domain with initial point in $T^S[p, q^s]$ iterates to a curve which crosses $T^S[q^u, r]$.

Lemma 7.17 *Let D be a regular domain for a quadrant Q such that $D \cap T^U = \partial D^U$ and D contains no smaller regular domains, and $D \cap T^P = \{p\}$. Then for any exact curve $\alpha : (I, \{0, 1\}) \rightarrow (\mathcal{C}^U T, T^S)$ with $\alpha(I) \subset D$, $\alpha(0) \in T^S[p, q^s]$ and $\alpha(1) \notin T^S[p, q^s]$, there exists k such that $f_{\min}^{nk}[\alpha]$ crosses $T^S[q^u, r]$.*

PROOF: Let S be the segment containing the final endpoint of α . Suppose $f^{in}(S) \subset D$ for all i . Then $f^{in}(S) \cap T^S[p, q^s] = \emptyset$ for all i , and since there are only finitely many stable segments in D , one must map into itself under f^{jn} for some j , giving a periodic point in D distinct from p , a contradiction. Therefore $f^{kn}(S) \not\subset D$ for some least k . For this k , we must have $f_{\min}^{kn}[\alpha] \cap T^S[q^u, r] \neq \emptyset$, since crossing $T^S[p, q^s]$ would imply a point of $f^{kn} \circ \alpha$ in $T^S(q^s, f^{-n}(q^s))$, a contradiction. $\square$

Recall the meaning of a curve $\tilde{\alpha}$ tightening onto α given in Definition 2.35. We show that if D and $\tilde{D}$ are regular domains for a quadrant, then some iterate of α_D tightens onto $\alpha_{\tilde{D}}$. The importance of this result is that if a curve $\hat{\alpha}$ tightens onto α , and both curves are tight curves embedded in a graph, then $\alpha(I) \subset \hat{\alpha}(I)$.

Lemma 7.18 *Let D be a regular domain for a quadrant Q such that $D \cap T^U = \partial D^U$, and let $\tilde{D} \subset D$ be a regular domain. Then there exists k such that $f_{\min}^{kn}[\alpha_D]$ tightens onto $[\alpha_{\tilde{D}}]$.*

PROOF: Let k be the least integer such that $f^{kn}(\tilde{q}^u) \notin T^U[p, q^u]$. Then $f^{kn}(\tilde{q}^u) \notin D$, so the endpoint of $f_{\min}^{kn}[\alpha_{\tilde{D}}]$ does not lie in D . Further, the first intersection of $f_{\min}^{kn}[\alpha_{\tilde{D}}]$ with ∂D^S must be with $T^S[q^u, r]$, so $f_{\min}^{kn}[\alpha_{\tilde{D}}]$ tightens onto $[\alpha_D]$, which is the curve from $T^S[p, q^s]$ to $T^S[q^u, r]$ in D with minimal intersections with T^S . $\square$

An example of the minimal iterates of a curve $\alpha = \alpha_D$ for a regular domain D is shown in Figure 52.

Lemma 7.19 *Let D be a regular domain for a quadrant Q , and $\tilde{D} \subset D$ a regular domain for Q such that $\tilde{D} \cap T^U = \partial \tilde{D}^U$. Let α be a curve in D such that $\alpha(0) \in T^S[p, q^s]$ and $\alpha(1) \in T^S[q^u, r]$. Then there exists k such that $f_{\min}^{kn}[\alpha]$ tightens onto $[\alpha_{\tilde{D}}]$.*




Figure 52: A curve α_D in a regular domain D and some of its minimal iterates.

PROOF: For any i , $f_{\min}^{in}[\alpha]$ must cross $T^S[q^u, r]$, and do so before it crosses $T^S[p, q^s]$. Choose k such that $f_{\min}^{kn}(\alpha(0)) \in T^S[p, \tilde{q}^s]$. Then $f_{\min}^{kn}[\alpha]$ tightens onto $[\alpha_{\tilde{D}}]$. $\square$

If Q is a quadrant with a regular domain D such that $D \cap T^U = D^U$, we let $D(Q)$ be the smallest regular domain for Q , and write $[\alpha_Q]$ for $[\alpha_{D(Q)}]$. Since for any regular domain D , the n th iterate of α_D tightens onto itself, there is some recurrent behaviour for the graph representative. If such a curve than has some iterate which tightens onto α_D for a different quadrant, the graph representative maps edges from one quadrant to another. This provides the definition of an alpha-chain.

Definition 7.20 (Alpha chain) Let $([f]; T)$ be a well-formed trellis mapping class and Q_U and Q_S be quadrants of T which are contained in regular domains which do not intersect T^U in their interiors. We say there is an *alpha chain* from Q_U to Q_S if there exists n such that $f_{\min}^n[\alpha_{Q_U}]$ tightens onto $[\alpha_{Q_S}]$.




Figure 53: A trellis with an alpha chain from Q_U to Q_S .

An example of an alpha chain from a quadrant Q_U to Q_S is shown in Figure 53. If there is an alpha chain from any segment to any other, the trellis is said to be *transitive*.

Definition 7.21 (Transitive trellis) We say that T is *transitive* if every quadrant Q is contained in a regular domain regular D_Q such that $D_Q \cap T^U = \partial D_Q$, and for every pair of quadrants Q_U and Q_S , there is an alpha chain from Q_U to Q_S .

The following lemma gives a sufficient condition for the existence of an alpha-chain from Q_U to Q_S .

Lemma 7.22 *Suppose D_U and D_S are regular domains for Q_U and Q_S respectively such that $D_U \cap D^S$ is a rectangular domain with unstable edges contained in ∂D_U^U and stable edges in ∂D_S^S . Then there is an alpha chain from Q_U to Q_S .*

PROOF: Let β be a curve in $D^U \cap D^S$ homotopic to the initial interval of $T^U(p_u) \cap D^S$. Then $f_{\min}^{k_u n_u}[\alpha_{Q_U}]$ tightens onto β for some k_u , and since β is a curve joining the stable sides of D_S , we also have $f_{\min}^{k_s n_s}[\beta]$ tightens onto α_{Q_S} for some k_s . Hence $f_{\min}^{k_u n_u + k_s n_s}[\alpha_{Q_U}]$ tightens onto $[\alpha_{Q_S}]$ as required. $\square$

Transitivity is preserved by taking stable extensions.

Lemma 7.23 *If there is an alpha chain from Q_U to Q_S for $([f]; T)$ and $(l\hat{f}; \hat{T})$ is a stable extension of T , then there is an alpha chain from Q_U to Q_S for $([\hat{f}]; \hat{T})$. In particular, if $([f]; T)$ is transitive, and $(l\hat{f}; \hat{T})$ is a stable extension of T , then $([\hat{f}]; \hat{T})$ is transitive.*

PROOF: By Lemma 7.18, $\hat{f}_{\min}^{k_u n_u}[\hat{\alpha}_{Q_U}]$ tightens onto $[\alpha_{Q_U}]$ for some k_u , and by Lemma 7.19, $\hat{f}_{\min}^{k_s n_s}[\alpha_{Q_S}]$ tightens onto $[\alpha_{Q_S}]$ for some k_s . Further, $f_{\min}^n[\alpha_{Q_U}]$ tightens onto $[\alpha_{Q_S}]$ for some n since there is an alpha-chain from Q_U to Q_S in T . Therefore, there is an alpha chain from Q_U to Q_S in $\tilde{T}$. $\square$

We now show that a transitive trellis has a transitive graph representative.

Lemma 7.24 *Let $([f]; T)$ be an transitive trellis mapping class. Then the graph representative $(g; G, W)$ of $([f]; T)$ has a single transitive component with positive topological entropy.*

PROOF: Let $\bar{G} = \bigcup_{n=0}^{\infty} g^n(G)$ be the essential subgraph, and let e be any expanding edge of G . By irreducibility, there is a quadrant Q such that $g^n(e)$ tightens to α_Q for some n .

We now consider preimages of edges. Again, let e be an expanding edge of G . There exists a tight curve β_0 in $\tilde{G}$ with endpoints in W such that β_0 is homotopic to a subinterval of a branch $T^U(p, b)$. We then find a curve β_1 in $\tilde{G}$ homotopic to a subinterval of a branch at $f^{-1}(p)$. Proceeding recursively gives an edge-path $\beta_n = \alpha_Q$ where β_i is a sub-path of $g(\beta_{i+1})$ for $0 \leq i < n$. Therefore, there exists a quadrant Q and an integer n such that $g^n(\alpha_Q) \supset e$.

Now if Q_U and Q_S are any two quadrants, there exists n such that $g^n(\alpha_{Q_U})$ contains a sub-curve α_{Q_S} . Since $(f; T)$ is irreducible, we can find N such that for any two quadrants and any $n \geq N$, the iterate $g^n(\alpha_{Q_U})$ contains a sub-path α_{Q_S} . Therefore, any such curve α generates the same graph component under iteration.

Combining these results shows that there exists N such that if e_1 and e_2 are any two edges, and $n \geq N$, then $e_2 \subset g^n(e_1)$. $\square$

Finally, we show that the hyperbolicity near T^P is enough to create intersections from which we can deduce regularity and transitivity. Note that for this result we are concerned with f -extensions of a trellis map $(f; T)$ rather than an isotopy class.

Lemma 7.25 *Let $(f; T)$ be a trellis map and Q a quadrant of T . Let $q \in T^U(Q)$ be the endpoint of an unstable segment S on the Q -side of $T^S(Q)$, and let $q^s \in T^S(Q)$ be the endpoint of a stable segment U on the Q -side of $T^S(Q)$. Then there exists k such that $f^{-kn}(S)$ intersects U at a point r such that $\{p, f^{-kn}(q), q^s, r\}$ are the vertices of a regular domain for Q .*

PROOF: By the Lambda lemma, as $i \rightarrow \infty$, $f^{-in}(S)$ limits on $W^S(Q)$ in the C^1 topology. Take a neighbourhood K of q^s such that $f^{-n}(K) \cap T^S[p, q^s] = \emptyset$, and choose k such that $f^{-kn}(S)$ intersects U in at a point r in K such that $T^S[q^s, r] \subset K$, and the domain D with vertices at $\{p, f^{-kn}(q), q^s, r\}$ is a rectangle which does not intersect $f^{-n}(K)$, and such that $f^n(T^U(p, f^{-kn}(q^u))) \cap D = T^U[p, f^{-kn}(q^u)]$. Let $\hat{T} = (T^U, f^{-nk}(T^S))$. Then $f^{-n}(T^U[q^s, r]) \cap D \subset f^{-n}(K) \cap D = \emptyset$, so D is a regular domain of $(f; \hat{T})$, as shown in Figure 54. $\square$




Figure 54: Backward iterates of S eventually form the opposite stable side of a regular domain.

7.3 Existence of entropy minimisers

To prove the existence of a diffeomorphism in a trellis mapping class whose topological entropy is the Nielsen entropy of the class, we first show how to construct such a diffeomorphism for a particularly simple class of trellises.

Lemma 7.26 *Let $([f]; T)$ be a trellis mapping class. Suppose every chaotic region R of $([f]; T)$ is a rectangle, and that every such rectangle is contained in a rectangular domain D with $D^U \subset f^{-1}(T^U)$ such that every region in D and $f(D)$ is a rectangle. Then there exists a diffeomorphism $\hat{f} \in ([f]; T)$ such that $h_{top}(\hat{f}) = h_{top}([f]; T)$.*

PROOF: For every domain D containing a chaotic region R , foliate D and $f(D)$ by an unstable foliation $\mathcal{F}^U$ parallel to T^U and a transverse stable foliation $\mathcal{F}^S$ parallel to T^S . Isotope f to obtain a diffeomorphism $\tilde{f}$ which preserves the stable and unstable foliations, and for which all points of non-chaotic regions are in the basin of a stable or unstable periodic orbit. Let (G, W) be a graph representative of $([f]; T)$ for which G is transverse to T^S , and $\pi : (\mathcal{C}^U T, T^S) \rightarrow (G, W)$ be a deformation-retract which collapses each leaf of $\mathcal{F}^S$ onto a point of g . Then $\pi \circ \tilde{f} = g \circ \pi$ on every chaotic region R , so $h_{top}(\tilde{f}) = h_{top}(g) = h_{niel}[f; T]$. $\square$

We can use this to prove the general case.

Theorem 7.27 (Existence of entropy minimisers) *Let $([f]; T)$ be a trellis mapping class. Suppose there is a diffeomorphism $\hat{f} \in ([f]; T)$ such that every extension of T by $\hat{f}$ is minimal. Then there is a uniformly-hyperbolic diffeomorphism $\tilde{f} \in ([f]; T)$ such that every extension of T by $\tilde{f}$ is minimal, and $h_{top}(\tilde{f}) = h_{niel}[f; T]$.*

PROOF: By Theorem 7.5, any trellis mapping class of the form $[\hat{f}; (T^U, \hat{f}^{-n}(T^S))]$ has the same Nielsen entropy at $([f]; T)$. By Lemma 7.25 we can therefore take a $\hat{f}$ -extension $\hat{T}_1$ of T such that every quadrant of $\hat{T}_1$ lies in a rectangular region, and by irreducibility, we can ensure that every region of $\hat{T}_1$ is a topological disc or annulus. If r_0 is a point of $\hat{T}_1^S$ at the end of a trivial branch and lies in a chaotic region, by introducing new unstable curves as in Theorem 7.11, we can take a minimal supertrellis $([\tilde{f}_2]; \tilde{T}_2)$ such that the orbit of r_0 lies in a repelling or chaotic region. Similarly, if a_0 is a point of $\tilde{T}_2^S$ at the end of a trivial branch of $\hat{T}_1^U$, by introducing new stable curves as in Theorem 7.11, we can take a minimal supertrellis $([\tilde{f}_3]; \tilde{T}_3)$ such that the orbit of a_0 lies in an attracting or chaotic region.

Now suppose there is a chaotic region R of $([\tilde{f}_3]; \tilde{T}_3)$ which is not a rectangle. Then the graph representative $(\tilde{g}_3; \tilde{G}_3, \tilde{W}_3)$ of $([\tilde{f}_3]; \tilde{T}_3)$ has a peripheral loop or a valence- n vertex in R which corresponds to a boundary component or essential periodic orbit which does not shadow T^S . Introducing new stable curves for all such R as in Theorem 7.11 gives a minimal supertrellis $([\tilde{f}_4]; \tilde{T}_4)$ for which R is a domain containing an attractor and some chaotic rectangles.

Every chaotic region R of $([\tilde{f}_4]; \tilde{T}_4)$ is now a rectangle. Taking a minimal backwards stable iterate $([\tilde{f}_5]; \tilde{T}_5)$ of $([\tilde{f}_4]; \tilde{T}_4)$ gives a trellis mapping class satisfying the conditions of Lemma 7.26. Hence there is a diffeomorphism $\tilde{f} \in ([\tilde{f}_5]; \tilde{T}_5)$ for which $h_{top}(\tilde{f}) = h_{niel}[\tilde{f}_5; \tilde{T}_5]$, which equals $h_{niel}[f; T]$ since every extension and supertrellis is minimal. Further, it is clear that every $\tilde{f}$ -extension of $\tilde{T}_5$ is minimal, so every $\tilde{f}$ -extension of T is minimal. $\square$

7.4 Approximate entropy minimising diffeomorphisms

Throughout this section we take $([f]; T)$ to be a well-formed irreducible trellis mapping class, $\lambda = \exp(h_{niel}[f; T])$ and $\lambda_\epsilon = \exp(h_{niel}[f; T] + \epsilon)$. We aim to show that we can always find a diffeomorphism $\hat{f} \in ([f]; T)$ which has entropy less than λ_ϵ . We first consider how to perform non-minimal extensions without increasing the Nielsen entropy above $h_{niel}[f; T] + \epsilon$. The technical details are given in Lemma 7.28 and Lemma 7.29, which deal directly with the graph representatives. We then prove Lemma 7.30, which allows us to consider non-minimal extensions without reference to the graph representative.

Our aim is to construct a trellis mapping class which satisfies the conditions of Theorem 7.27. To do this, we may need to introduce new periodic points to the trellis to create attracting and repelling regions. We then iterate curves bounding an attractor or repeller

into regular domains, and finally take non-minimal iterates to move bigon boundaries into attractors and repellers.




Figure 55: A trellis and its graph representative in the neighbourhood of a periodic control edge z_p .

To show that we can take non-minimal iterates without increasing the Nielsen entropy, we consider a length function on a graph near a periodic control edge, as shown in Figure 55. We let x^u be the first intersection of $T^U(Q)$ with T^S , and $S_X(Q)$ be the stable segment on the Q -side of $T^U(Q)$ and x^u , as shown. Let z_p be the control edge crossing $S_P(Q)$, and z_x the control edge crossing $S_X(Q)$. We let $\beta(Q)$ be the curve from the periodic control vertex v_p to the periodic control vertex v_x . Note that α is homotopic to the inclusion of the path $z_p\beta z_x$ in M .

Note that for any graph representative of a trellis mapping class, we can choose a length function l such that $l(z) = l(p) = l_p$ for all control edges z and all peripheral edges p , and such that $l(g(e)) < \lambda_\epsilon l(e)$ for all edges e .

Lemma 7.28 *Let $([f_i]; T_i)$ be the minimal extension of $([f]; T)$ with $([f_0]; T_0) = ([f]; T)$ and $T_i = (T^U, f_i^{-1}(T_{n-1}))$. Let $(g_i; G_i, W_i)$ be the graph representative of $([f_i]; T_i)$ with natural embeddings $(G_{i-1}; W_{i-1}) \hookrightarrow (G_i, W_i)$, and let l_i be a length function on $(G_i; W_i)$ preserved by this embedding. Let Q be a quadrant of $([f]; T)$, let z_p be the control edge crossing $S_P(Q)$, and let z_i be the control edge crossing the first stable segment crossing $T^U(Q)$. Let β_i be the edge-path in G_i with initial vertex at v_p and final vertex at v_i in z_i . Then $l_i(\beta_i) \rightarrow 0$ as $i \rightarrow \infty$, and $l(z_i) \rightarrow 0$ as $i \rightarrow \infty$.*

PROOF: The result follows since $g_i(\beta_i) = \beta_{i-1}$, and β_i does not contain any periodic control edges. $\square$

Lemma 7.29 *Let Q_0 is a quadrant with $f(Q_0) = Q_1$ and $2l(\beta(Q_1)) + 2l(z_x(Q_1)) < (\lambda_\epsilon - 1)l_p$. Then there is a well-formed trellis mapping class $([\hat{f}]; \hat{T})$ such that $h_{niel}([\hat{f}]; \hat{T}) < \lambda_\epsilon$ and $U_X(Q)$ intersects $\hat{f}^{-1}(S_X(f(Q)))$.*

PROOF: To obtain the graph $\hat{G}$, subdivide the control edge $z_p(Q_0)$ into two control edges $\hat{z}_p$ and $\hat{z}_q$, and two expanding edges, one edge $\hat{e}_p$ joining $\hat{p}$ to $\hat{q}$, and one edge $\hat{e}_q$ joining $\hat{q}$ to $v_p(Q_0)$ as shown in Figure 56.

We then recursively subdivide control edges mapping to z_p into a control edge eventually mapping to $\hat{z}_p$ and an expanding edge. A compatible graph map is obtained by taking $\hat{e}_p$ to $\beta(Q_1)$, $\hat{z}_q$ to $z_x(Q_1)$, and $\hat{e}_q$ to $\bar{z}_x(Q_1)\bar{\beta}(\bar{Q}_1)$. Since this graph map takes $z_p(Q_0)$, with length l_p , to $z_p(Q_1)\beta(Q_1)z_x(Q_1)\bar{z}_x(Q_1)\bar{\beta}(\bar{Q}_1)$ with length less than $\lambda_\epsilon l_p$, but is unchanged




Figure 56: A non-minimal extension.

on all other edges, the growth rate is still less than λ_ϵ . Hence the graph representative of $([\hat{f}]; \hat{T})$, which can be obtained by Algorithm 5.7, has growth rate less than λ_ϵ . $\square$

We now use these technical results to show how to create intersections bounding regular domains.

Lemma 7.30 *Let $([f]; T)$ be a trellis mapping class and Q a quadrant of $([f]; T)$. Let $q^u \in T^U(Q)$ be the endpoint of an unstable segment S_0 on the same side of $T^S(Q)$ as Q , and let $q^s \in T^S(Q)$ be the endpoint of a stable segment U on the same side of $T^S(Q)$ as Q . Further, for any $\epsilon > 0$ there is a minimal stable extension $([\hat{f}]; \hat{T})$ and an integer k such that $\hat{f}^{-kn}(S) \subset \hat{T}^S$ and intersects U at a point r such that $\{p, f^{-kn}(q^u), q^s, r\}$ are the vertices of a regular domain for Q .*

PROOF: Let S_i be the segment of $f^{-k}(S_0)$ containing $f^{-in}(q^u)$, let z_i be the control edge crossing S_i . Let z_p be the control edge crossing $S_P(Q)$, and let β_i be the edge path from z_p to z_i which contains z_i as its final edge, but does not contain z_p . Then the hypotheses of the length functions for the graph representative imply that $l(\beta_i) \rightarrow 0$ as $k \rightarrow \infty$. Fix k such that $1 + 2l(\beta_k) < \lambda_\epsilon$. $\square$

Before proving the main theorem we show how to ensure that every quadrant Q is contained in a regular domain whose opposite sides bound attractors or repellers.

Lemma 7.31 *Let $([f]; T)$ be a transitive trellis mapping class, and P be an essential periodic orbit of $([f]; T)$ which does not shadow T^P . Then there is a minimal supertrellis $([\tilde{f}]; \tilde{T})$ of $([f]; T)$ obtained by blowing up at P such that every quadrant Q is contained in a regular rectangular region with the opposite sides not in T .*

PROOF: Since P is an essential periodic orbit, every curve $\alpha(Q)$ eventually maps across P , so crosses a stable segment of P . Hence iterating this segment backwards gives a curve bounding a regular domain, and further backward iterates give a regular region. A similar argument holds for forward iterates of unstable segments, since in this case we can simply reverse time. $\square$

Theorem 7.32 (Existence of approximate entropy minimisers) *Let $([f]; T)$ be a well-formed trellis mapping class. Then for every $\epsilon > 0$, there exists a diffeomorphism $\hat{f} \in ([f]; T)$ such that $h_{top}(\hat{f}) < h_{niel}[f; T] + \epsilon$.*

PROOF: We repeatedly construct supertrellises $([f_i]; T_i)$ with $h_{niel}[f_i; T_i] < h_{niel}[f; T] + \epsilon = \log \lambda_\epsilon$.

Take $([f_0]; T_0) = ([f]; T)$. By irreducibility, there is a minimal extension $([f_1]; T_1)$ of $([f_0]; T_0)$ such that for every pair of nontrivial branches $T_1^U(p_u, b_u)$ and $T_1^S(p_s, b_s)$, there are points $p_u = p_0, p_1, \dots, p_n = p_s$ such that $T_1^U(p_u, b_u) \cap T_1^S(p_1) \neq \emptyset$, $T_1^U(p_i) \cap T_1^S(p_{i+1}) \neq \emptyset$ for $1 \leq i < n-1$ and $T_1^U(p_{n-1}) \cap T_1^S(p_s, b_s) \neq \emptyset$. By Lemma 7.30, there is a non-minimal extension $([f_2]; T_2)$ of $([f_1]; T_1)$ such that every quadrant of T_2 is contained in a regular rectangular region.

We next construct a non-minimal extension $([f_3]; T_3)$ of $([f_2]; T_2)$ such that every nontrivial branch of T_3^U intersects every nontrivial branch of T_3^S . If $T^U(Q_U)$ intersects $T^S(Q)$ and $T^U(Q)$ intersects $T^S(Q_S)$, we can take minimal iterates of $T^U(Q_U)$ and $T^S(Q_S)$ until $T^U(Q_U) \supset U_X(Q)$ and $T^S(Q_S) \supset S_X(Q)$. Then an application of Lemma 7.30 shows that there is a non-minimal intersection such that $T^U(Q_U) \cap T^S(Q_S) \neq \emptyset$.

Now construct a non-minimal extension $([f_4]; T_4)$ of $([f_3]; T_3)$ such that there is an alpha-chain from any quadrant Q_U of T_4 to any other quadrant Q_S . Since $T_3^S(Q_S)$ intersects $T_3^U(Q_U)$, by taking minimal backward iterates of $T_3^S(Q_S)$ we can ensure that $S_X(Q_U) \subset T^S(Q_S)$. Then by Lemma 7.30, there is a non-minimal extension such that $T^S(Q_S)$ crosses a regular region $R(Q_U)$. Similarly, we can ensure $T^U(Q_U)$ crosses the regular region $R(Q_S)$. A further application of Lemma 7.30 to $S_X(Q_U)$ gives a non-minimal extension with an alpha-chain from Q_U to Q_S . Repeating this construction for all regions gives the required trellis T_4 .

Let $([f_5]; T_5)$ be a minimal supertrellis of $([f_4]; T_4)$ such that all trivial branches of T_4 are contained in a non-chaotic region, as given by Theorem 7.11. If $([f_5]; T_5)$ has no attractors or no repellers, take a further minimal supertrellis $([f_6]; T_6)$, with at least on attractor and one repeller, as given by Theorem 7.11 and Theorem 7.11. Since $([f_6]; T_6)$ is transitive, we can take a minimal extension $([f_7]; T_7)$ by iterating the stable and unstable curves bounding stable and unstable regions such that all quadrants of T_7 are contained in a regular region with the opposite stable side bounding an attracting domain, and the opposite unstable side bounding a repelling domain.




Figure 57: Backward iterates of B give an inner bigon in a repelling domain.

Now, for any inner bigon B such that B^S is nonwandering, we take backward iterates of B^S to obtain a stable segment S in a regular domain of a quadrant Q . Applying Lemma 7.30 gives a non-minimal extension such that $f^{-n}(B)$ contains a single inner bigon, and this inner bigon lies in a repelling domain, as shown in Figure 57. Similarly, by taking forward iterates of B^U we can ensure $f^n(B)$ contains a single inner bigon in an attracting domain. Applying this procedure to all bigons gives a trellis mapping class $([f_8]; T_8)$.

Finally, we take a minimal supertrellis $([f_9]; T_9)$ satisfying the conditions of Lemma 7.26 by introducing new stable curves as in Theorem 7.11. Then there is a uniformly-hyperbolic diffeomorphism $\hat{f} \in ([f_9]; T_9)$, and hence in $([f]; T)$ such that $h_{top}(\hat{f}) = h_{niel}[f_9; T_9] < h_{niel}[f; T] + \epsilon$ as required. $\square$

We now prove the existence of pseudo-Anosov map which approximate the Nielsen entropy. The condition on attractors and repellers is to ensure that the trellis mapping class contains a pseudo-Anosov map, since pseudo-Anosov maps have no attractors or repellers. The strategy is to create the periodic orbits which will give the one-prong singularities of the pseudo-Anosov map. Many of the steps of the proof mimic those the proof of Theorem 7.32.

Theorem 7.33 (Existence of pseudo-Anosov representatives) *Let $([f]; T)$ be a trellis mapping class with no attractors or repellers. Then for any $\epsilon > 0$ there exists a pseudo-Anosov diffeomorphism $\hat{f} \in ([f]; T)$ such that $h_{top}(\hat{f}) < h_{top}([f]; T) + \epsilon$.*

PROOF: If T has an adjacent pair of trivial branches, take an extension $([f_1]; T_1)$ such that these two branches intersect in a single transverse homoclinic point. This does not affect the Nielsen entropy. If T has any other trivial branches, take a minimal extension $([f_2]; T_2)$ such that these branches have an intersection point, which is possible by Theorem 7.11 since T has no attracting or repelling regions. As in the proof of Theorem 7.32, take an extension $([f_3]; T_3)$ which is transitive. Take a further extension $([f_4]; T_4)$ such that every branch intersects every other with both orientations, which is possible by transitivity since we can take minimal iterates such that every unstable branch has an extension with and iterate of every stable branch which does not map to an intersection of T_3 , and an isotopy in the neighbourhood of this intersection yields three intersections, one with the opposite orientation.




Figure 58: The domain with vertices q_0 and q_1 must have periodic orbits in the rectangles R_0 and R_1 .

Let B be an inner bigon of T_4 , as shown in Figure 58(a). By Lemma 7.30, we can find a stable extension such that $f^{-n}(B^S)$ crosses a regular domain $R(Q_U)$ for some n . Further, by removing intersections if necessary, we can ensure that $f^{-n}(B^S)$ crosses D twice and gives a new inner bigon $\hat{B}$, as shown in Figure 58(b). A further application of Lemma 7.30 shows that we can find a stable extension $[\hat{f}_3; \hat{T}_3]$ such that $\hat{B}^U$ crosses some regular domain, as shown in Figure 58(c). Since the regions R are mapped over by $\alpha(Q_U)$ and map over $\alpha(Q_S)$, they must contain a periodic orbit, since T is transitive. Applying

this construction for every inner bigon of $([f_4]; T_4)$ gives a non-minimal extension $([f_5]; T_5)$ such that every bigon of T_4 is a domain of T_5 containing an essential periodic orbit of $([f_5]; T_5)$.

Let $([f_6]; T_6)$ be the trellis mapping class obtained by puncturing at a periodic orbit in every inner bigon of T^4 to give a surface M_6 . Then $h_{niel}[f_6; T_6] = h_{niel}[f_5; T_5]$. Since T_4 is a subtrellis of T_6 , we can take a trellis mapping class $([f_7]; T_7) = ([f_6]; T_4)$ in the surface M_6 . Since every inner bigon of T_4 contains component of ∂M_6 , the trellis mapping class $([f_7]; T_7)$ has no inner bigons. The graph representative $(g_7; G_7, W_7)$ is locally injective except at cusps, so is efficient and hence is a train-track map for a pseudo-Anosov diffeomorphism $\tilde{f} \in ([f_7]; T_7)$. Then $h_{top}(\tilde{f}) = h_{niel}[f_7; T_7] \leq h_{niel}[f_6; T_6] < h_{niel}[f; T] + \epsilon$ as required. $\square$

7.5 Non-existence of entropy minimisers

Suppose that an irreducible trellis mapping class has a uniformly hyperbolic diffeomorphism realising the Nielsen entropy. The following lemma shows that any isotopy removing intersections results in a trellis mapping class with strictly smaller Nielsen entropy.

Lemma 7.34 *Let $([f]; T)$ be a well-formed irreducible trellis mapping class, and suppose f be a uniformly-hyperbolic diffeomorphism such that $h_{top}(f) = h_{niel}[f; T]$. Then if $([\bar{f}]; \bar{T})$ is a trellis mapping class which does not force $([f]; T)$, then $h_{niel}[\bar{f}; \bar{T}] < h_{niel}[f; T]$.*

PROOF: By taking an f -extension T_1 of T , we can ensure that every inner bigon of




Figure 59: Change in trellis and graph at a homoclinic bifurcation.

$([f]; T_1)$ is contained in a larger domain with the topology of Figure 59(a), and that these domains are separate. Further, we can ensure that every quadrant is contained in a rectangular region. Now consider the effect of a homotopy to a trellis mapping class $([f_2]; T_2)$ removing one pair of orbits on the same inner bigons. The resulting trellis and graph locally have the topology of the right of Figure 59(b). The edges a_0 and a_2 must be expanding, since they forward iterate to a closed segment, and since every quadrant is contained in a rectangular region, $\partial g^n(a_0) = \partial g^n(a_2) = b$ for some edge b . Therefore a_0 and a_2 are folded together to obtain the graph representative for $([f_2]; T_2)$. Further, there must be an edge mapping $\dots \bar{a}_0 a_2 \dots$. Hence the entropy of the graph representative of $([f_2]; T_2)$ is less than that of $[f; T]$, so $h_{niel}[f_2; T_2] < h_{niel}[f; T]$. Since we can then prune $([f_2]; T_2)$ to obtain $([\bar{f}]; \bar{T})$, we have $h_{niel}[\bar{f}; \bar{T}] \leq h_{niel}[f_2; T_2]$, hence $h_{niel}[\bar{f}; \bar{T}] < h_{niel}[f; T]$ as required. $\square$

We use this to prove that the sufficient condition given for the existence of entropy minimisers in Theorem 7.27 is necessary.

Theorem 7.35 (Non-existence of entropy minimisers) *Let $([f]; T)$ be an irreducible trellis mapping class. Suppose that for every diffeomorphism $\hat{f} \in ([f]; T)$ there is a $\hat{f}$ -extension of T which is not minimal. Then there does not exist a diffeomorphism in $([f]; T)$ whose topological entropy equals $h_{\text{niel}}[f; T]$.*

PROOF: Suppose there is an f -extension T_1 of T which is not minimal. Let T_2 be a transitive f -extension of T_1 , and T_3 be a further extension such that every non-wandering segment of T crosses a regular domain of T_3 . Then we have entropy bound $h_{\text{top}}(f) \geq h_{\text{niel}}[f; T_3]$.

Take $([f_4]; T_4)$ to be a minimal supertrellis of $([f]; T_3)$ for which the opposite sides of every regular region bound a stable or unstable region. Then every nonwandering segment of T enters a stable or unstable region of T_4 . Prune $([f_4]; T_4)$ to obtain a trellis mapping class $([f_5]; T_5)$ which forces $([f]; T_1)$ satisfying the conditions of Theorem 7.27. Then $h_{\text{niel}}[f_3; T_3] = h_{\text{niel}}[f_4; T_4] \geq h_{\text{niel}}[f_5; T_5]$, but by Lemma 7.34, $h_{\text{niel}}([f_5]; T_5) > h_{\text{niel}}[f; T]$. Combining these inequalities we have $h_{\text{top}}(f) \geq h_{\text{niel}}[f_5; T_5] > h_{\text{niel}}[f; T]$ as required. $\square$

The results of this section show that many fundamental properties of an irreducible trellis type $[f; T]$ depends on whether there is a diffeomorphism $f \in [f; T]$ for which $h_{\text{top}}(\tilde{f}) = h_{\text{niel}}[f; T]$. If such a diffeomorphism exists, then the entropy of the trellis type is carried in a uniformly hyperbolic diffeomorphism, but is fragile in the sense that any pruning will reduce the Nielsen entropy, and any diffeomorphism $\hat{f}$ for which some extension is non-minimal must have strictly greater topological entropy. If no such diffeomorphism exists, every diffeomorphism in the trellis type has an extension which is non-minimal, but pruning this extension gives a trellis type for which the entropy is still greater than the Nielsen entropy of $[f; T]$.

A Intersections and Isotopies of Curves in Surfaces

In this section we prove some technical results on isotopies of curves in surface. (A *curve* is a function $\alpha : I \rightarrow M$; a *path* is the image of such a function.) We consider a surface M with a set $\Gamma = \{\gamma_1 \dots \gamma_k\}$ of finitely-many cross-cuts. (A *cross-cut* is a simple curve or path with both endpoints but no interior points in ∂M .) All homotopies and isotopies of cross-cuts will be assumed to fix endpoints. Homotopies and isotopies of curves with endpoints on cross-cuts will be assumed to keep this property.

The main results are Theorem A.6, which gives conditions under which two sets of curves can be isotoped to reduce the number of intersections with each other, Theorem A.8, which shows how to isotope curves in a controlled way, and Theorem A.11, which shows that there is essentially a unique configuration of curves with minimal intersections in any isotopy class.

We begin by quoting some standard results of (surface) topology. The first is due to Epstein [Eps66], the can be found in [Hir76], and the third is an easy consequence of

the classification of orientable surfaces. All these results also hold in the differentiable category.

Theorem A.1 *Let α_0 and α_1 be two simple curves in a surface M with the same endpoints. Then α_0 and α_1 are homotopic if and only if they are isotopic (relative to endpoints).*

Theorem A.2 *Suppose α_0 and α_1 are two simple curves in a surface M which are isotopic via an isotopy α_t . Then there is an isotopy $h_t : M \rightarrow M$ with $h_0 = \text{id}$ such that $h_t(\alpha_0) = \alpha_s$.*

Theorem A.3 *Let M be a surface and Γ a cross-cut in M . Then if α_0 and α_1 are curves in $M \setminus \Gamma$ which are homotopic in M , then α_0 and α_1 are homotopic in $M \setminus \Gamma$. An alternative statement is that the inclusion-induced map $\pi_1(M \setminus \Gamma) \rightarrow \pi_1(M)$ is injective.*

We also have the following differentiable version of the Alexander trick:

Theorem A.4 *Suppose M is a polygon, and $f : M \rightarrow M$ is a smooth map which preserves the vertices and edges of M . Then f is isotopic to the identity.*

A.1 Minimal intersections

A particularly important concept will be that of two sets of cross-cuts having *minimal intersections* with each other. We give a definition based on a local definition, and then show that this implies that the two sets have minimise the total number of intersections in the isotopy class.

Definition A.5 (Minimal intersections) A set of mutually disjoint cross-cuts $\aleph$ has *minimal intersections* with Γ if $\aleph$ and Γ are transverse and there is no disc in M bounded by an arc in $\aleph$ and an arc in Γ .




Figure 60: Intersections of a curve with two cross-cuts. (a) Not minimal intersections. (b) An isotopic curve with minimal intersections.

The following elementary result relates this local definition to the global property of having fewest intersections in the isotopy class.

Theorem A.6 *Let M be a surface, Γ a set of mutually disjoint cross-cuts in M , and α a cross-cut. Then there is an isotopy h_t such that $h_0 = \text{id}$ and $h_1 \circ \alpha$ has minimal intersections with Γ . Further, $|(h_1 \circ \alpha) \cap \Gamma| \leq |\alpha \cap \Gamma|$, with equality if and only if α has minimal intersections with Γ .*

The proof is an easy consequence of the following lemma:

Lemma A.7 *Suppose α is a curve in M , and D a disk bounded by an arc in Γ and an arc of α , and that the intersection of α and γ is (topologically) transverse. Let U be a simply-connected neighbourhood of D such that $\alpha \cap \Gamma \cap U \subset D$. Then there is an isotopy $h_t : M \rightarrow M$ supported in U such that $h_1 \circ \alpha$ and α agree outside of U , but $h_1 \circ \alpha$ has no intersections with Γ in U .*

PROOF: Let $\alpha[t_1, t_2]$ be the arc of α bounding D , and t_0, t_3 be parameters such that $\alpha[t_0, t_1)$ and $\alpha(t_2, t_3]$ lie in $U \setminus \Gamma$. Then there is a smooth curve α which agrees with α except in (t_0, t_2) , and $\alpha \cap \Gamma \cap U = \emptyset$. There is a homeomorphism h_1 of M supported in U such that $h_1 \circ \alpha = \alpha$, and since U is simply-connected, h_1 is isotopic to id by an isotopy supported in U . $\square$

PROOF OF THEOREM A.6: If the intersections of α with Γ are not isolated, we take a small perturbation h_{t_0} which is isotopic to the identity and for which $h_{t_0} \circ \alpha$ has isolated intersections with Γ . In particular, α_{t_0} has finitely many intersections with Γ while α has infinitely many. Now, if α_{t_0} does not have minimal intersections with Γ , there is a disc D bounded by an arc in Γ and an arc of α_{t_0} , so by Lemmas A.7, we can isotope further by h_t to obtain $\alpha_{t_1} = h_{t_1} \circ \alpha$ with two fewer intersections with $h_{t_0} \circ \alpha$. (If one of the bounding points of D is an endpoint of α , we can still isotope to remove one intersection.) We continue isotoping until we reach a curve α_1 with minimal intersections with Γ . The procedure must terminate in a finite number of steps, since α_{t_0} has finitely many intersections. $\square$

A.2 Ambient isotopies of curves

Theorem A.8 *Let M be a surface and Γ a set of cross-cuts in M . Then if h is isotopic to id and $h(\Gamma) = \Gamma$ there is an isotopy h_t such that $h_0 = \text{id}$, $h_1 = h$ and $h_t(\Gamma) = \Gamma$.*

Lemma A.9 *Suppose α_0 and α_1 are homotopic curves which do not intersect Γ . Then there is an isotopy h_t such that $h_0 = \text{id}$, $h_1(\alpha_0) = \alpha_1$ and $h_t|_{\Gamma} = \text{id}$.*

PROOF: By Theorem A.3, Then $\hat{\alpha}_t = \pi \circ \tilde{\alpha}$ is a homotopy between α_0 and α_1 in $M \setminus \Gamma$, and hence in a subsurface K of M such that $K \cap \Gamma = \emptyset$. Then by Theorem A.1, there is an isotopy α_t between α_0 and α_1 supported in K . By Theorem A.2, there is an isotopy h_t supported in K such that $h_1(\alpha_0) = \alpha_1$. $\square$

This result can be extended to collections of curves.

Lemma A.10 *Let $\aleph_0 = \{\alpha_0^1 \dots \alpha_0^n\}$ and $\aleph_1 = \{\alpha_1^1 \dots \alpha_1^n\}$ be two sets of disjoint curves in a surface M such that α_0^i is homotopic to α_1^i for $i = 1, \dots, n$. Then there is an isotopy h_t such that $h_0 = id$ and $h_1(\alpha_0^i) = \alpha_1^i$.*

PROOF: Assuming the result for $n-1$ curves, we can find an isotopy such that $h_t(\alpha_0^i) = \alpha_1^i$ for $i \leq n-1$, and $h_1(\alpha_0^n)$ and α_1^n are both disjoint from $\alpha_1^1 \dots \alpha_1^{n-1}$ and isotopic in M . The result follows by taking $\Gamma = \{\alpha_1^1, \dots, \alpha_1^{n-1}\}$ in Lemma A.9. $\square$

We can now prove Theorem A.8.

PROOF: Let $\aleph$ be a set of cross-cuts which is disjoint from Γ such that $M \setminus (\aleph \cup \Gamma)$ is simply-connected. By Lemma A.10, there is an isotopy $\tilde{h}_t$ such that $\tilde{h}_0 = id$, $\tilde{h}_1(\aleph) = h(\aleph)$ and $\tilde{h}_t(\Gamma) = \Gamma$. Then $\tilde{h}_1^{-1} \circ h(\Gamma) = \Gamma$ and $\tilde{h}_1^{-1} \circ h(\aleph) = \aleph$, so $\tilde{h}_1^{-1} \circ h$ is a homeomorphism/diffeomorphism of a polygon, and clearly must preserve the sides of the polygon. Hence by Theorem A.4, there is an isotopy $\hat{h}_t$ such that $\hat{h}_0 = id$, $\hat{h}_1 = \tilde{h}_1^{-1} \circ h$ and $\hat{h}_t(\Gamma) = \Gamma$. Let $h_t = \tilde{h}_t \circ \hat{h}_t$. Then $h_0 = id$, $h_1 = h$ and $h_t(\Gamma) = \tilde{h}_t + t(\hat{h}_t(\Gamma)) = \tilde{h}_t(\Gamma) = \Gamma$ as required. $\square$

A.3 Nielsen intersection theory

Theorem A.11 *Let M be a surface, Γ a set of cross-cuts in M . Then if $\aleph_0$ and $\aleph_1$ are homotopic sets of cross-cuts with minimal intersections with Γ , there is a isotopy h_t such that $h_0 = id$, $h_1(\aleph_0) = \aleph_1$ and $h_t(\Gamma) = \Gamma$. In particular, h_1 is a homeomorphism $(M, \aleph_0, \Gamma) \longrightarrow (M, \aleph_1, \Gamma)$.*

To prove Theorem A.11, we need to develop a form of Nielsen theory for intersections of curves on surfaces. We first define an *intersection class*, and then show that each class has a well-defined *intersection number*.

Definition A.12 (Intersection class) Let Γ be a set of cross-cuts in a surface M , α a curve in M , and x_0 and x_1 points in $\alpha \cap \Gamma$, with $x_i = \alpha(s_i)$ for $i = 0, 1$. We say x_0 and x_1 lie in the same *intersection class* if $\alpha|_{[s_0, s_1]}$ is homotopic to a curve in Γ .

If α_t is an isotopy, and there are functions $\sigma, \tau : I \longrightarrow I$ such that $\alpha_{\tau(t)}(\sigma(t)) \in \Gamma$ for all t , then we say $\alpha_{\tau(0)}(\sigma(0))$ and $\alpha_{\tau(1)}(\sigma(1))$ are *connected* by the isotopy; essentially they lie in equivalent intersection classes.

Lemma A.13 *Suppose α_t is an isotopy of curves, and there are functions σ and τ such that $\alpha_{\tau(0)}(\sigma(0))$ and $\alpha_{\tau(1)}(\sigma(1))$ are connected by the isotopy α_t , and $\tau(0) = \tau(1) = \tau$. Then $\alpha_\tau(\sigma(0))$ and $\alpha_\tau(\sigma(1))$ lie in the same intersection class.*

PROOF: Let

$$\hat{\alpha}_t(s) = \alpha_{(1-t)\tau + t\tau(s)}((1-t)((1-s)\sigma(0) + \sigma(1)) + t\sigma(s)).$$

Then $\hat{\alpha}_0(s) = \alpha_\tau((1-s)\sigma(0) + \sigma(1))$, $\hat{\alpha}_1(s) = \alpha_{\tau(s)}(\sigma(s))$ and $\hat{\alpha}_t(i) = \alpha_\tau(\sigma(i))$ for $i = 0, 1$. $\square$

Lemma A.14 Suppose α_t is a homotopy, and there are functions $\sigma_i, \tau_i : I \longrightarrow I$ for $i = 0, 1$ such that $\alpha_{\tau_i(t)}(\sigma_i(t)) \in \Gamma$ for $i = 1, 2$. Then the curves $\alpha_i|_{[\sigma_i(0), \sigma_i(1)]}$ are homotopic via curves with endpoints in Γ .

PROOF: A homotopy is given by

$$\hat{\alpha}_t(s) = \alpha_{(1-s)\tau_0(t)+s\tau_1(t)}((1-s)\sigma_0(t) + s\sigma_1(t)),$$

since we can check that $\hat{\alpha}_i(s) = \alpha_i((1-s)\sigma_0(i) + s\sigma_1(i)) = \alpha_i|_{[\sigma_i(0), \sigma_i(1)]}(s)$ and $\hat{\alpha}_t(i) = \alpha_{\tau_i(t)}(\sigma_i(t)) \in \Gamma$. $\square$

It is clear that for a given curve, the intersection classes are open in the intersection set.

Definition A.15 (Intersection number) The *intersection number* of an intersection class is the topological intersection number of that class. An intersection class is *essential* if its intersection number is non-zero.

We now show that essential intersection classes preserve their relative ordering under homotopies of curves.

Lemma A.16 Let α_0 and α_1 be isotopic curves in M with endpoints in Γ and minimal intersections with Γ . Let t_0^k and t_1^k be ordered parameter values of the intersections of α_0 and α_1 with Γ respectively. Then for each curve γ_i , there are integers $k_1 \dots k_{n_i}$ such that the intersections of α_0 and α_1 with γ_i are $t_0^{k_1} \dots t_0^{k_{n_i}}$ and $t_1^{k_1} \dots t_1^{k_{n_i}}$ respectively.

PROOF: Since α_0 and α_1 have minimal intersections with Γ , any two intersections of α_0 or α_1 with Γ are transverse and lie in different intersection classes. Let α_t be an isotopy between α_0 and α_1 , let $X = \{s, t : \alpha_t(s) \in \Gamma\}$, and let $X_t = \{s : \alpha_t(s) \in \Gamma\}$. Since points in the same component of X with the same t -value must correspond to points in the same intersection class of α_t , for each intersection point of α_0 there is a connected subset of X which contains a point of X_t for any t ; in particular, it contains a point of X_1 , and this point must be unique since α_1 also has minimal intersections with Γ . This gives the required bijection from $\alpha_0 \cap \Gamma$ to $\alpha_1 \cap \Gamma$. $\square$

Lemma A.17 Let α_0 and α_1 be isotopic curves in M with endpoints in Γ and minimal intersections with Γ . Parameterise α_0 and α_1 so that the intersections occur at the same parameter values $t_1, t_2, \dots, t_n$. Then there are tubular neighbourhoods U, V of Γ with $\text{cl}(U) \subset V$, and a homeomorphism supported on V , such that $h(\alpha_0)$ and α_1 agree on U , and consist of curves, each crossing Γ once transversely. Further, on each interval $[t_i, t_{i+1}]$ for $i < n$ and for the intervals $[0, t_1]$ and $[t_n, 1]$, the curves $h \circ \alpha_0$ and α_1 are homotopic.

PROOF: By Lemma A.16, we can find a U, V and h such that $h(\alpha_0)$ and α_1 agree on U , so need only show that $h(\alpha_0)$ and α_1 are consist of mutually homotopic curves outside U . By Lemma A.14, the curves are homotopic on each interval. $\square$

Using these results, we can prove Theorem A.11.

PROOF: By Lemma A.16 we can find an isotopy h_t such that $h_1(\alpha_0)$ and α_1 agree on a neighbourhood U of Γ . Without loss of generality, therefore, we can assume that α_0 and α_1 agree on U . Now let V be a tubular neighbourhood of Γ such that $\text{cl}(V) \subset U$, and K be the surface $M \setminus V$. Then $\alpha_0 \cap K$ and $\alpha_1 \cap K$, give families of mutually disjoint homotopic curves, and the result follows from Lemma A.10. $\square$

References

- [BH92] Mladen Bestvina and Michael Handel. Train tracks and automorphisms of free groups. *Ann. of Math. (2)*, 135(1):1–51, 1992.
- [BH95] Mladen Bestvina and Michael Handel. Train-tracks for surface homeomorphisms. *Topology*, 34(1):109–140, 1995.
- [Bir49] Garrett Birkhoff. Theorie et applications des treillis. *Ann. Inst. H. Poincaré*, 11:227–240, 1949.
- [BW95] Keith Burns and Howard Weiss. A geometric criterion for positive topological entropy. *Comm. Math. Phys.*, 172(1):95–118, 1995.
- [CB88] A. Casson and S. Bleiler. *Automorphisms of surfaces, after Nielsen and Thurston*, volume 9 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, UK, 1988.
- [Col99] Pieter Collins. Dynamics forced by surface trellises. In Marcy Barge and Krystyna Kuperberg, editors, *Geometry and Topology in Dynamics*, number 246 in Contemporary Mathematics. American Mathematical Society, Providence, 1999.
- [Col01] Pieter Collins. Relative periodic point theory. *Topology Appl.*, 115(1):97–114, 2001.
- [Con78] Charles Conley. *Isolated Invariant Sets and the Morse Index*, volume 38 of *CBMS Regional Conference Series in Mathematics*. American Mathematical Society, Providence, 1978.
- [CP96] Freddy Christiansen and Antonio Politi. Symbolic encoding in symplectic maps. *Nonlinearity*, 9(6):1623, 1996.
- [dC99] André de Carvalho. Pruning fronts and the formation of horseshoes. *Ergodic Theory Dynamical Systems*, 19(4):851–894, 1999.
- [dCH] André de Carvalho and Toby Hall. The forcing relation for horseshoe braid types. Preprint.
- [Eas86] Robert Easton. Trellises formed by stable and unstable manifolds in the plane. *Trans. Amer. Math. Soc.*, 294(2):719–732, 1986.

- [Eas98] Robert W. Easton. *Geometric methods for discrete dynamical systems*. Number 50 in Oxford Engineering Science Series. Oxford University Press, Oxford, 1998.
- [Eps66] David Epstein. Curves on 2-manifolds and isotopies. *Acta Math.*, 115:83–107, 1966.
- [FM93] John Franks and Michael Misiurewicz. Cycles for disk homeomorphisms and thick trees. In *Nielsen Theory and Dynamical Systems*, number 152 in Contemporary Mathematics, pages 69–139, 1993.
- [Fri83] David Fried. Periodic points and twisted coefficients. 1007:261–293, 1983.
- [Han85] Michael Handel. Global shadowing of pseudo-Anosov homeomorphisms. *Ergodic Theory Dynamical Systems*, 5(3):373–377, 1985.
- [Han99] Michael Handel. A fixed-point theorem for planar homeomorphisms. *Topology*, 38(2):235–264, 1999.
- [Hir76] Morris W. Hirsch. *Differential Topology*. Number 33 in Graduate Texts In Mathematics. Springer-Verlag, Berlin, 1976.
- [Hul00] Helen Hulme. *PhD. Thesis*. PhD thesis, University of Liverpool, 2000.
- [Jia93] Boju Jiang. Nielsen theory for periodic orbits and applications to dynamical systems. In Christopher McCord, editor, *Nielsen Theory and Dynamical Systems*, number 152 in Contemporary Mathematics, pages 183–202. American Mathematical Society, Providence, 1993.
- [Kei97] Ulrich Keil. *Construction Algorithmique D’un Collage Efficace Pour Les Homéomorphismes D’une Surface À Bord (Algorithmic Construction of an Efficient Splicing for Homeomorphisms of a Punctured Surface)*. PhD thesis, L’Université Paul Sabatier De Toulouse, 1997.
- [KO98] Bernd Krauskopf and Hinke Osinga. Growing 1d and quasi-2d unstable manifolds of maps. *J. Comput. Phys.*, 146(1):404–419, 1998.
- [Los93] Jérôme Los. Pseudo-Anosov maps and invariant train tracks in the disc: A finite algorithm. *Proc. London Math. Soc. (3)*, 66(2):400–430, 1993.
- [Los96] Jérôme Los. On the conjugacy problem for automorphisms of free groups. *Topology*, 35(3):779–806, 1996.
- [MT94] F. Allan McRobie and J. M. T. Thompson. Knot-types and bifurcations sequences of homoclinic and transient orbits of a single-degree-of-freedom driven oscillator. *Dynam. Stab. Sys.*, 9:223–251, 1994.
- [MV93] L. Mora and M. Viana. Abundance of strange attractors. *Acta Math.*, 171:1–71, 1993.

- [New80] Sheldon E. Newhouse. Lectures on dynamical systems. In *Dynamical Systems (CIME Summer School, Bressanone, 1978)*, volume 8 of *Progress in Mathematics*, pages 1–114. Birkhäuser, 1980.
- [Poi99] Henri Poincaré. *Les méthodes nouvelle de la mécanique céleste*. 1892–1899.
- [Ray98] Victoria Rayskin. Phd thesis, university of california, berkeley. Ph.D. Thesis, 1998.
- [RK94] Vered Rom-Kedar. Homoclinic tangles—classification and applications. *Nonlinearity*, 7(2):441–473, 1994.
- [SDM99] D. Sterling, Holgar R. Dullin, and J.D. Meiss. Homoclinic bifurcations for the hénon map. *Phys. D*, 134:153–184, 1999.
- [Sim89] C. Simó. On the analytical and numerical approximation of invariant manifolds. In D. Benest and C. Froeschlé, editors, *Les Méthodes Modernes de la Mécanique Céleste*, pages 285–329. Goutelas, 1989.
- [Sma63] Stephen Smale. A structurally stable differentiable homeomorphism with an infinite number of periodic points. In *Proceedings of the International Symposium on Nonlinear Vibrations*, volume 2, pages 365–366, 1963.